

THE GROWTH RATE OF SYMPLECTIC HOMOLOGY AND AFFINE VARIETIES

MARK MCLEAN

ABSTRACT. We will show that the cotangent bundle of a manifold whose free loop space homology grows exponentially is not symplectomorphic to any smooth affine variety. We will also show that the unit cotangent bundle of such a manifold is not Stein fillable by a Stein domain whose completion is symplectomorphic to a smooth affine variety. For instance, these results hold for end connect sums of simply connected manifolds whose cohomology with coefficients in some field has at least two generators. We use an invariant called the growth rate of symplectic homology to prove this result.

CONTENTS

1. Introduction	1
2. Main definitions and properties	6
2.1. Liouville domains	6
2.2. Symplectic homology	8
2.3. Growth rates	12
3. Growth rate linear algebra	13
4. Growth rate geometry	15
4.1. Some alternate definitions of growth rate	15
4.2. Growth rate of cotangent bundles	36
5. Compactifications of algebraic varieties	41
5.1. Making the divisors orthogonal	41
5.2. Making the smooth affine variety nice at infinity	51
6. The growth rate of affine varieties	60
7. Finite covers of smooth affine varieties	69
8. Appendix A: convex symplectic manifolds	70
9. Appendix B: A Maximum Principle	75
References	77

1. INTRODUCTION

The aim of this paper is to use an invariant called the growth rate of symplectic homology to construct many examples of cotangent bundles which are not symplectomorphic to any smooth affine variety. We also have a contact analogue of this result which says that if we take the unit cotangent

bundle of one of these manifolds then it is not Stein fillable by a Stein domain whose completion is symplectomorphic smooth affine variety. All of our manifolds are assumed to be oriented unless explicitly stated otherwise.

If we have some Liouville domain M (defined in Section 2.1) and a class $b \in H^2(M, \mathbb{Z}/2\mathbb{Z})$, then the growth rate of symplectic homology is an invariant $\Gamma(M, b) \in \{-\infty\} \cup [0, \infty)$. This invariant was originally defined in [Sei08, Section 4]. Every Liouville domain also has a completion $\widehat{M}$, and the growth rate is in fact an invariant of the completion $\widehat{M}$ up to symplectomorphisms preserving the class b and so we will often write $\Gamma(\widehat{M}, b)$.

A smooth affine variety A has a symplectic structure obtained by embedding it into $\mathbb{C}^N$ algebraically and then pulling back the standard symplectic form onto A . It turns out that this is symplectomorphic to the completion of some Liouville domain $\overline{A}$ obtained from taking a large closed ball in $\mathbb{C}^N$ and intersecting it with A . Because the symplectic form is a biholomorphic invariant of A , we can assign the invariant $\Gamma(A, b) = \Gamma(\overline{A}, b)$ (see [EG91]).

We say that a contact manifold C is *algebraically Stein fillable* if it is Stein fillable by a Stein domain D whose completion $\widehat{D}$ is symplectomorphic to a smooth affine variety A . For a smooth affine variety A we also have a finite invariant $m_A \in \mathbb{N}_{>0}$ defined as follows: choose a compactification X of A by a smooth normal crossing divisor (i.e. $X \setminus A$ is a union of smooth transversely intersecting connected complex hypersurfaces S_i , $i = 1, \dots, k$). For $I \subset \{1, \dots, k\}$ we write $S_I := \cap_{i \in I} S_i$. Let $d := \max\{n - \dim_{\mathbb{C}}(S_I) \mid S_I \neq \emptyset\}$ where $n = \dim_{\mathbb{C}} A$. We define m_A to be the minimum of d over all compactifications described above.

The main result of this paper is:

Theorem 1.1. $\Gamma(A, b) \leq m_A$ for any class $b \in H^2(A, \mathbb{Z}/2\mathbb{Z})$. In particular $\Gamma(A, b) < \infty$.

In fact we will prove a more general theorem:

Theorem 1.2. Suppose that the boundary of a Liouville domain M is algebraically Stein fillable by A , then $\Gamma(M, b) \leq m_A$ for any $b \in H^2(M, \mathbb{Z}/2\mathbb{Z})$.

This theorem will be proven in Section 6. These theorems have been proven in [Sei08, Section 4] when A has complex dimension 2. Let Q be a compact manifold. Choose some metric on Q and let $\mathcal{L}^q Q$ be the space of all free loops on Q of length less than or equal to q , and let $\mathcal{L}Q$ be the space of all free loops on Q . Let $\mathbb{K}$ be any field and let $a^{\mathbb{K}}(q)$ be the rank of the image of the inclusion map

$$H_*(\mathcal{L}^q(Q), \mathbb{K}) \hookrightarrow H_*(\mathcal{L}(Q), \mathbb{K}).$$

The manifold Q is said to have *exponential growth* if for some $\mathbb{K}$, the function $a^{\mathbb{K}}(q)$ grows faster than any polynomial (we mean here that for any polynomial $p(q)$, $a^{\mathbb{K}}(q) > p(q)$ for q large enough). If the manifold Q has finite fundamental group, then this definition of exponential growth is the same as saying that $\sum_{i=1}^k b_i(\mathcal{L}(Q), \mathbb{K})$ grows faster than any polynomial in k for

some $\mathbb{K}$ by [Gro78]. There are many examples of manifolds with exponential growth. Here are some:

- (1) Any surface of genus 2 or higher.
- (2) The end connect sum of two simply connected manifolds $M_1 \# M_2$ where $H^*(M_1, \mathbb{K})$ has at least two generators and $H^*(M_2, \mathbb{K})$ has rank at least 3 (See [Lam01]).
- (3) Any manifold Q whose fundamental group is the free product of at least 3 non-trivial groups (See Lemma 4.20, and Lemma 4.21).

Conjecturally (see [VP84]) there should be many more manifolds with exponential growth such as simply connected manifolds whose Betti numbers are greater than that of the torus. Given any Riemannian manifold Q , we can construct a symplectic manifold T^*Q where locally the symplectic form is $\sum_i dq_i \wedge dp_i$ where q_i are the position coordinates and p_i are the momentum coordinates. We also have a Liouville domain D^*Q which is the manifold with boundary consisting of covectors of length less than or equal to 1. Its boundary is the contact manifold denoted by S^*Q consisting of covectors of length 1 and the contact form here is $\sum_i p_i dq_i$.

By using work from [SW06, Corollary 1.2] (although we could have used ideas from [AS06] or [Vit96]), it can be shown that $\Gamma(T^*Q, \omega_2) = \infty$ when Q has exponential growth and ω_2 is the pullback of the second Stiefel-Whitney class of Q . Hence we have the following corollary of Theorem 1.1:

Corollary 1.3. *If Q has exponential growth then T^*Q is not symplectomorphic to any smooth affine variety.*

This is because $\Gamma(T^*Q, \omega_2) = \infty$ and $\Gamma(A, b) < \infty$ for any smooth affine variety A and any $b \in H^2(A, \mathbb{Z}/2\mathbb{Z})$. We also have the following corollary of Theorem 1.2:

Corollary 1.4. *The unit cotangent bundle of Q is not algebraically Stein fillable.*

If we wish to consider unoriented manifolds Q then we need to do the following: A contact manifold C is said to be *covered algebraically Stein fillable* if there is a Stein filling M of C whose completion $\widehat{M}$ is symplectomorphic to a finite cover of a smooth affine variety.

Theorem 1.5. *Suppose the boundary of a Liouville domain M is covered algebraically Stein fillable by a smooth affine variety A , then $\Gamma(M, b) \leq m_A$ for any $b \in H^2(M, \mathbb{Z}/2\mathbb{Z})$.*

We will prove this theorem in Section 7. We then immediately get the following corollaries:

Corollary 1.6. *If Q is a compact manifold of exponential growth which is possibly unoriented such that some finite cover is oriented and is has exponential growth then T^*Q is not symplectomorphic to any smooth affine variety.*

This corollary is true for the following reason: If T^*Q is symplectomorphic to some smooth affine variety A then for every oriented finite cover $\tilde{Q}$ of Q , $T^*\tilde{Q}$ is symplectomorphic to some finite cover of A . Hence by the above theorem we have that $\Gamma(T^*\tilde{Q}, \omega_2) < \infty$ which means that $\tilde{Q}$ cannot have exponential growth.

Similar reasoning also gives us the following corollary:

Corollary 1.7. *The unit cotangent bundle of any (possibly unoriented) Q of exponential growth with dimension > 2 is not covered algebraically Stein fillable. In particular the unit cotangent bundle of any such Q is not algebraically Stein fillable.*

The reason why we need dimension > 2 is that if we have some algebraic Stein filling C of S^*Q then we require that $\pi_1(Q) = \pi_1(S^*Q) = \pi_1(C)$ so that we can take an appropriate cover of C . In future work the author hopes to prove a based loop space version of the above theorems using wrapped Floer homology. This will enlarge the class of manifolds whose cotangent bundle is not symplectomorphic to a smooth affine variety. For instance this would be true if their Betti numbers are greater than that of the torus.

There is a similar conjecture relating smooth affine varieties with cotangent bundles which we will now describe. Suppose that U is a real affine algebraic variety over $\mathbb{R}$ which is diffeomorphic to our manifold Q . We will suppose that Q is simply connected. We say that Q has a *good complexification* if there exists a U as described above such that the natural map $U(\mathbb{R}) \rightarrow U(\mathbb{C})$ is a homotopy equivalence. There is a question in [Tot03] which asks if Q has a Riemannian metric of non-negative sectional curvature when Q has a good complexification. There is also a conjecture attributed to Bott which conjectures that any simply connected manifold with non-negative sectional curvature is rationally elliptic. Rationally elliptic means that $\dim \pi_*(Q) \otimes \mathbb{Q} < \infty$ (see [FHT82]). This conjecture is mentioned in [FHT01, Question 12, page 519]. Felix and Thomas in [FT82] proved that if a simply connected manifold is rationally elliptic then the growth rate of the Betti numbers of its based loop space grows sub exponentially. Vigué-Poirrier in [VP84, Page 415] conjectured that the growth rate of the Betti numbers of the based loop space grow exponentially if and only if the same thing is true for the free loop space. If we look at all of these statements then we get the following question: Suppose Q is simply connected of exponential growth, then is it true that Q does not admit a good complexification? This question has some similarities with Corollary 1.3 because we can choose a symplectic form on $U(\mathbb{C})$ such that a neighbourhood of $U(\mathbb{R})$ is symplectomorphic to a neighbourhood of the zero section of T^*Q and also because both T^*Q and $U(\mathbb{C})$ deformation retract onto Q .

We will now give a brief sketch of the proof of Theorem 1.2. Symplectic homology of $\widehat{M}$ is the homology of a chain complex having the following generators:

- (1) Critical points of some Morse function on M .

(2) Two copies of each closed Reeb orbit of ∂M .

This chain complex has a natural filtration by $\mathbb{R}_{\geq 0}$ called the *action*. Critical points have action 0 and the action of each Reeb orbit is its length. Hence we can define $SH_*^{\leq \lambda}(M)$ for each λ as the homology of the subcomplex of orbits of action $\leq \lambda$. The growth rate is the rate at which

$$\text{rank im}(SH_*^{\leq \lambda}(M) \rightarrow \varinjlim_l SH_*^{\leq l}(M))$$

grows with respect to λ . For instance if the rank is bounded above by some polynomial of degree k then the growth rate is less than or equal to k . It turns out that our smooth affine variety A is symplectomorphic to $\widehat{M}$ for some M . We need to provide an upper bound for the growth rate of M . One way of doing this is to show that the number of Reeb orbits of length $\leq \lambda$ is bounded above by some polynomial of degree at most m_A . This is the method used in [Sei08, Section 4] when Theorem 1.2 is proven in complex dimension 2. This upper bound is achieved by compactifying A by a smooth normal crossing divisor and carefully constructing a Liouville domain using this divisor. The problem is that this upper bound is much harder to achieve in higher dimensions. In this paper we define growth rate in a slightly more flexible way so that it is easier to provide such an upper bound.

We have some Hamiltonian $H \geq 0$ and an almost complex structure J called a *growth rate admissible pair* which satisfies certain technical properties. From this pair (H, J) we can define a series of homology groups $SH_*^\#(\lambda H, J)$ generated by 1-periodic orbits of λH . For $\lambda_1 \leq \lambda_2$ there is a natural map

$$SH_*^\#(\lambda_1 H, J) \rightarrow SH_*^\#(\lambda_2 H, J).$$

We prove that the growth rate of

$$\text{rank im}(SH_*^\#(\lambda H, J) \rightarrow \varinjlim_l SH_*^\#(lH, J))$$

is equal to the original definition of growth rate (Section 4.1). Finally we carefully construct a Hamiltonian H so that there is a degree m_A polynomial bound on the number of 1-periodic orbits of a very small perturbation of λH and this gives us an upper bound on the growth rate (Section 6). We construct this Hamiltonian by first compactifying our smooth affine variety A with a smooth normal crossing divisor D . We deform D without changing the symplectic structure on its complement A so that there is a nice symplectic structure near D (Section 5). We can construct H so that around some point $p \in D$ we have that D looks like $\{x_1 = 0, \dots, x_k = 0, y_1 = 0, \dots, y_k = 0\}$ where $x_1, \dots, x_n, y_1, \dots, y_n$ is some nice coordinate chart around p and H is a product Hamiltonian. The symplectic structure is not quite the standard one on this chart but the Hamiltonian flow of H is exactly the same as the Hamiltonian flow with respect to the standard symplectic structure. This enables us to find all the

1-periodic orbits of H and give a bound proportional to $\lambda^k \leq \lambda^{m_A}$ in each chart. Hence we get an upper bound of m_A .

Acknowledgements: I would like to thank Mohammed Abouzaid, Dietmar Salamon, Burt Totaro, Paul Seidel and Ivan Smith for extremely useful comments. In addition, the author is grateful to the referee for many valuable comments and suggestions. The author was partially supported by NSF grant DMS-1005365.

2. MAIN DEFINITIONS AND PROPERTIES

2.1. Liouville domains. A *Liouville domain* is a compact manifold N with boundary and a 1-form θ_N satisfying:

- (1) $\omega_N := d\theta_N$ is a symplectic form.
- (2) The ω_N -dual of θ_N is transverse to ∂N and pointing outwards.

We will write X_{θ_N} for the ω_N -dual of θ_N . Sometimes we have manifolds with corners with 1-forms θ satisfying the same properties as above. We view these as Liouville domains by smoothing the corners slightly. By flowing ∂N backwards along X_{θ_N} we have a small collar neighbourhood of ∂N diffeomorphic to $(0, 1] \times \partial N$ with $\theta_N = r_N \alpha_N$. Here r_N parameterizes $(0, 1]$ and α_N is the contact form on the boundary given by $\theta_N|_{\partial N}$. The completion $\hat{N}$ is obtained by gluing $[1, \infty) \times \partial N$ to this collar neighbourhood and extending θ_N by $r_N \alpha_N$.

The Liouville domains that we will be studying are called Stein domains. A *Stein manifold* is a complex manifold that can be properly embedded in $\mathbb{C}^N$. An equivalent definition of a Stein manifold is a complex manifold admitting an exhausting plurisubharmonic function. A function is *exhausting* if it is bounded from below and the preimage of every compact set is compact. A function $f : S \rightarrow \mathbb{R}$ is *plurisubharmonic* if $(-dd^c f)(X, JX) > 0$ for all non-zero vectors X where $d^c := d \circ J$ and J is the complex structure on the complex manifold S . This implies that $\omega_f := -dd^c f$ is a symplectic form. If c is a regular value of an exhausting plurisubharmonic function f then $f^{-1}(-\infty, c]$ is a Liouville domain with Liouville form $-d^c f$. We call such a domain a *Stein domain*. A Stein manifold is of *finite type* if it admits an exhausting plurisubharmonic function with only finitely many singularities. A plurisubharmonic function f is said to be *complete* if the ω_f -dual of $-d^c f$ is integrable. Every finite type Stein manifold admits an exhausting plurisubharmonic function which is complete. Also any two such functions on S give symplectomorphic Stein manifolds. Hence the symplectic form is a biholomorphic invariant. We will write ω_S for this symplectic form. If f is finite type then it turns out for $C \gg 0$ that (S, ω_S) is symplectomorphic to the completion of $f^{-1}(-\infty, C]$. One way of obtaining such a symplectic form is embedding S as a closed proper holomorphic embedding into $\mathbb{C}^N$ and then pulling back the standard symplectic form ω_{std} on $\mathbb{C}^N$. Important examples of finite type Stein manifolds A are smooth affine varieties (see [Sei08, Section 4b]). In fact from [Sei08, Section 4b] we have for any smooth

affine variety A , a Liouville domain $\bar{A}$ which is unique up to isotopy through Liouville domains such that $\widehat{\bar{A}}$ is symplectomorphic to A . Uniqueness here means that if B is isomorphic as a variety to A then $\bar{A}$ is isotopic to $\bar{B}$. From now on throughout this paper if we have a smooth affine variety A we will write $\bar{A}$ for such a Liouville domain. We will call any such Liouville domain an *algebraic Stein domain*. Here is a direct way of constructing this Liouville domain: Choose any algebraic embedding ι of A into $\mathbb{C}^N$ (so it is a closed subvariety). Let (r_i, ϑ_i) be polar coordinates for the i 'th factor in $\mathbb{C}^N$. Let $R := \sum_i \frac{r_i^2}{4}$. We have $\theta_A := -d^c R = \sum_i \frac{r_i^2}{2} d\vartheta_i$. We have that $d\theta_A$ is equal to the standard symplectic structure on $\mathbb{C}^N$. By abuse of notation we write θ_A for $\iota^* \theta_A$, and $\omega_A := d\theta_A$.

Lemma 2.1. *There is a $C > 0$ such that for all $c \geq C, ((R|_A)^{-1}(-\infty, c], \theta_A)$ is a Liouville domain whose completion is symplectomorphic to (A, ω_A) .*

Proof. of Lemma 2.1. We will first show that for some $C \geq 0$, $R|_A$ has no singularities when $R|_A \geq C$. Compactify $\mathbb{C}^N$ to $\mathbb{P}^N$ and let X be the projective subvariety which is the closure of A in $\mathbb{P}^N$. Let H be the hypersurface $X \setminus A$. We have a line bundle $\mathcal{O}(1)$ described as follows: If we think of a point on $\mathbb{P}^N$ as a line Λ in $\mathbb{C}^{N+1}$ then the fiber at this point is the quotient $(\mathbb{C}^{N+1})^*/V$ where V is the vector subspace of linear 1-forms whose kernel contains Λ . The hyperplane H is represented by some linear hyperplane $\tilde{H}$ in $\mathbb{C}^{N+1}$ and so we have a section s given by a non-zero linear 1-form whose kernel is $\tilde{H}$. This section vanishes on H with order 1.

Another way of describing the fiber of this bundle over the point represented by the line Λ is as the space of linear 1-forms W_Λ which vanish in the hyperplane orthogonal to Λ . We can put a metric $\|\cdot\|$ on W_Λ induced by the standard metric on $(\mathbb{C}^{N+1})^*$ hence we view $\|\cdot\|$ as a metric on $\mathcal{O}(1)$. The curvature form is a positive $(1,1)$ form and this gives us a symplectic form on $\mathbb{P}^N$. We have an action of $U(N+1)$ on $\mathbb{P}^N$ induced by the standard action on $\mathbb{C}^{N+1}$. This action also naturally lifts to an action on the total space of $\mathcal{O}(1)$. The metric $\|\cdot\|$ is invariant under the action of $U(N+1)$. We can ensure that $U(N) \subset U(N+1)$ is the subgroup which sends H to H . This subgroup also fixes our section s so $u^*\|s\| = \|s\|$ for each $u \in U(N)$. Hence $\|s\|$ must be a function of R . If it wasn't then there exists two elements $a, b \in \mathbb{C}^N$ of the same modulus such that $\|s\|(a) \neq \|s\|(b)$ but this is impossible as there is a $u \in U(N)$ such that $u(a) = b$ which implies that $\|s\|(a) = u^*(\|s\|)(b) = \|s\|(b)$. The function $-\log \|s\|_A$ is a plurisubharmonic function on A . It is also equal to $f(R|_A)$ for some function f . We will first show that $-\log \|s\|_A$ has no singularities near infinity. If D was a smooth normal crossing divisor then [Sei08, Section 4b] tells us that $d \log \|s\|_A$ is non-zero near infinity. But the problem is that D could be anything. So by the Hironaka resolution of singularities [Hir64] we blow up X away from A to $\tilde{X}$ where $\tilde{X}$ is smooth and the pullback $\tilde{D}$ is a smooth normal crossing divisor. Let L be the pullback of $\mathcal{O}(1)$ and $\tilde{s}$ the pullback

of s . We can also pull back the metric $\|\cdot\|$ to L . To show that $d\log\|s\|$ is non-zero we just show that $d\log\|\tilde{s}\|$ is non-zero in exactly the same way as in Lemma [Sei08, Section 4b]: Let $p \in \tilde{D}$ and choose local holomorphic coordinates $z_1, \dots, z_n$ and a trivialization of L around p so that $s = z_1^{w_1} \cdots z_n^{w_n}$ ($w_i \geq 0$). The metric $\|\cdot\|$ on L is equal to $e^\psi |\cdot|$ for some function ψ with respect to this trivialization where $|\cdot|$ is the standard metric on $\mathbb{C}$. So

$$d\log\|\tilde{s}\| = -\psi - \left(\sum_i w_i d\log|z_i|\right).$$

If we take the vector field $Y := -z_1 \partial_{z_1} \cdots - z_n \partial_{z_n}$, then $d\log(|z_j|)(Y) = -1$ and $Y.\psi$ tends to zero. Hence $d\log\|\tilde{s}\|$ is non-zero near infinity which implies that $f(R) = -\log\|s\|$ has no singularities near infinity. Because $f(R)$ and R are both exhausting, we get that $f' > 0$ near infinity which implies that $R|_A$ has no singularities in the region $R \geq C$ for some $C > 0$.

Because A is a holomorphic submanifold of $\mathbb{C}^N$, we have that ω_A is a symplectic form. Let $c \geq C$ and write $A_c := (R|_A)^{-1}(-\infty, c]$. We have that A_c is a Liouville domain because if X_{θ_A} is the ω_A -dual of θ_A then it is equal to the gradient of $R|_A$ and hence is transverse to ∂A_c and pointing outwards because c is a regular value of $R|_A$ and $f'(c) > 0$. Also we have that the ω_A -dual X_{θ_A} of θ_A is an integrable vector field because $d(R|_A)(X_{\theta_A}) \leq dR(\sum_i \frac{r_i}{2} \frac{\partial}{\partial r_i})|_A = R|_A$ and hence if $\Phi_t^{X_{\theta_A}}$ is the flow of this vector field then $(R|_A)(\Phi_t^{X_{\theta_A}}(x))$ increases at a rate of at most $e^R|_A$. This first inequality is true because X_{θ_A} is the orthogonal projection of $\sum_i \frac{r_i}{2} \frac{\partial}{\partial r_i}$ onto TA so its length l decreases, but $R(\sum_i \frac{r_i}{2} \frac{\partial}{\partial r_i}) = \sup_{|V| \leq l} R(V)$. By flowing ∂A_c along X_{θ_A} we get that $\overline{A \setminus A_c}$ is diffeomorphic to $[1, \infty) \times \partial A_c$ and $\theta_A = r_A \alpha_{A_c}$ where r_A parameterizes $[1, \infty)$ (this is because $d(R|_A)(X_{\theta_A}) > 0$ for $R|_A \geq C$). Hence A is symplectomorphic to $\widehat{A_c}$. $\square$

A contact manifold C is said to be *Stein fillable* if it is contactomorphic to the boundary of some Stein domain. It is said to be *algebraically Stein fillable* if the completion of this Stein domain is symplectomorphic to A .

2.2. Symplectic homology. Let N be a Liouville domain with $c_1 = 0$. We make some additional choices $\eta := (\tau, b)$ for N . The element τ is a choice of trivialization of the canonical bundle of N up to homotopy and b is an element of $H^2(N, \mathbb{Z}/2\mathbb{Z})$. We will assume that ∂N has discrete period spectrum $\mathcal{P}_N$ (the set of lengths of Reeb orbits of the $(\partial N', \alpha_{N'})$). For each pair of numbers $a < b$ where $a, b \in [-\infty, \infty]$ we will define a symplectic homology group $SH_*^{(a,b)}(N, \eta)$ which is an invariant of (N, θ_N) up to exact symplectomorphism (although we suppress θ_N in the notation unless the context is unclear). Here an exact symplectomorphism between two Liouville domains N and N' is a smooth diffeomorphism Φ from N to N' such that $\Phi^* \theta_{N'} = \theta_N + df$ for some smooth function $f : N \rightarrow \mathbb{R}$.

A family of Hamiltonians $H : S^1 \times \widehat{N} \rightarrow \mathbb{R}$ is said to be *admissible* if $H(t, x) = \lambda r_N(x)$ near infinity where r_N is the cylindrical coordinate of $\widehat{N}$

and λ is some positive constant. Here λ is some positive constant which is not in the period spectrum of ∂N . We also require that $H_t|_N < 0$. We sometimes view H as a time dependent Hamiltonians $H_t : \widehat{N} \rightarrow \mathbb{R}$ where $t \in S^1$. We have an S^1 family of vector fields X_{H_t} and it has an associated flow $\Phi_{X_{H_t}}^t$ (a family of symplectomorphisms parameterized by $t \in \mathbb{R}$ satisfying $\frac{\partial}{\partial t} \Phi_{X_{H_t}}^t = X_{H_t}$ where we identify $S^1 = \mathbb{R}/\mathbb{Z}$). A 1-periodic orbit $o : S^1 \rightarrow \widehat{N}$ is a map which satisfies $o(t) = \Phi_{X_{H_t}}^t(x)$ for some $x \in \widehat{N}$. We say that o is non-degenerate if $D\Phi_{X_{H_t}}^1 : T_x \widehat{N} \rightarrow T_x \widehat{N}$ has no eigenvalue equal to 1. By [DS94], we can perturb H by a C^∞ small amount so that all of its 1-periodic orbits are non-degenerate. The problem here is that this Hamiltonian may not be admissible after perturbing it, so we need a lemma:

Lemma 2.2. *Let $H : S^1 \times N \rightarrow \mathbb{R}$ where N is a symplectic manifold. Let U be a small neighbourhood of some of the 1-periodic orbits of H such that no 1-periodic orbits intersect the boundary ∂U . Then we can perturb H by a C^∞ small amount to $\tilde{H}$ such that all the 1-periodic orbits of $\tilde{H}$ are non-degenerate inside U and $\tilde{H} = H$ outside U .*

Proof. of Lemma 2.2. We can choose U small enough so that the time 1 flow of X_{H_t} is well defined. Let U' be a smaller open neighbourhood of the 1-periodic orbits whose closure is contained in U . By slightly extending the work of [DS94], we can find for any positive function $f : N \rightarrow \mathbb{R}$, a Hamiltonian H_f such that $\|H_f - H\|_{C^\infty} < f$ and such that all the 1-periodic orbits of H_f are non-degenerate. Let R be an open neighbourhood of $\partial U'$ inside U where H has no 1-periodic orbits which intersect the closure of R . Choose a bump function $\rho : N \rightarrow \mathbb{R}$ which is 0 on a neighbourhood of $U' \setminus R$ and 1 on a neighbourhood of $N \setminus U'$. Let $\tilde{H}_f := \rho H + (1 - \rho)H_f$. Suppose for a contradiction that for every f sufficiently small, there exists a 1-periodic orbit of $\tilde{H}_f$ inside U which is degenerate. This orbit must intersect R because all orbits of H_f are non-degenerate away from R . Then by a compactness argument, we have a sequence of such orbits converging to a 1-periodic orbit of H which intersects the closure of R . This is impossible. Hence we have for any f arbitrarily small, there is a Hamiltonian H_f satisfying $\|H_f - H\|_{C^\infty} < f$ such that all of its orbits inside U are non-degenerate and $H_f = H$ outside U . $\square$

Because we have a trivialization τ of the canonical bundle of N , this gives us a canonical trivialization of the symplectic bundle TN restricted to an orbit o . Using this trivialization, we can define an index of o called the **Robbin-Salamon** index (This is equal to the Conley-Zehnder index taken with negative sign). We will write $i(o)$ for the index of this orbit o . For a 1-periodic orbit o we define the **action** $A_H(o)$:

$$A_H(o) := - \int_0^1 H(t, \gamma(t)) dt - \int_o \theta_N.$$

Choose a coefficient field $\mathbb{K}$. Let

$$CF_k^d(H, J, \eta) := \bigoplus_o \mathbb{K}\langle o \rangle$$

where we sum over 1-periodic orbits o of H satisfying $A_H(o) \leq d$ whose Robbin-Salamon index is k . We write

$$CF_k^{(c,d]}(H, J, \eta) := CF_k^d(H, J, \eta) / CF_k^c(H, J, \eta).$$

We need to define a differential for the chain complex $CF_k^d(H, J, \eta)$ such that the natural inclusion maps $CF_k^c(H, J, \eta) \hookrightarrow CF_k^d(H, J, \eta)$ for $c < d$ are chain maps. This makes $CF_k^{(c,d]}(H, J, \eta)$ into a chain complex as well. In order to define this we choose an $\mathbb{S}^1$ family of almost complex structures J_t compatible with the symplectic form. We assume that J_t is convex with respect to this cylindrical end outside some large compact set (i.e. $\theta \circ J_t = dr$). We also say that J_t is **admissible**.

We will now describe the differential

$$\partial : CF_k^d(H, J, \eta) \rightarrow CF_{k-1}^d(H, J, \eta).$$

We consider curves $u : \mathbb{R} \times \mathbb{S}^1 \rightarrow \widehat{N}$ satisfying the perturbed Cauchy-Riemann equations:

$$\partial_s u + J_t(u(s, t)) \partial_t u = \nabla^{g_t} H$$

where ∇^{g_t} is the gradient associated to the $\mathbb{S}^1$ family of metrics $g_t := \omega(\cdot, J_t(\cdot))$. For two periodic orbits o_-, o_+ let $\overline{U}(o_-, o_+)$ denote the set of all curves u satisfying the Cauchy-Riemann equations such that $u(s, \cdot)$ converges to $x_\pm$ as $s \rightarrow \pm\infty$. This has a natural $\mathbb{R}$ action given by translation in the s coordinate. Let $U(o_-, o_+)$ be equal to $\overline{U}(o_-, o_+)/\mathbb{R}$. For a C^∞ generic admissible complex structure we have that $U(o_-, o_+)$ is an $i^C(o_-, o_+) - 1$ dimensional oriented manifold (see [FHS95]). There is a maximum principle which ensures that all elements of $U(o_-, o_+)$ stay inside a compact set K (see [Oan04, Lemma 1.5], [AS10, Lemma 7.2] or Corollary 9.3). Hence we can use a compactness theorem (see for instance [BEH⁺03]) which ensures that if $i(o_-) - i(o_+) = 1$, then $U(o_-, o_+)$ is a compact zero dimensional manifold. The class $b \in H^2(N, \mathbb{Z}/2\mathbb{Z})$ enables us to orient this manifold (see [Abo10, Section 3.1]). Let $\#U(x_-, x_+)$ denote the number of positively oriented points of $U(x_-, x_+)$ minus the number of negatively oriented points. Then we have a differential:

$$\begin{aligned} \partial : CF_k^d(H, J, \eta) &\longrightarrow CF_{k-1}^d(H, J, \eta) \\ \partial\langle o_- \rangle &:= \sum_{i(o_-) - i(o_+) = 1} \#U(o_-, o_+) \langle o_+ \rangle \end{aligned}$$

By analyzing the structure of 1-dimensional moduli spaces, one shows $\partial^2 = 0$ and defines $SH_*(H, J, \eta)$ as the homology of the above chain complex. As a $\mathbb{K}$ vector space $CF_k^d(H, J, \eta)$ is independent of J and b , but its boundary operator does depend on J . The homology group $SH_*^d(H, J, \eta)$ depends on

H and η but is independent of J up to canonical isomorphism. We define $SH_*^{(c,d]}(H, J, \eta)$ as the homology of the chain complex

$$CF_*^d(H, J, \eta)/CF_*^c(H, J, \eta).$$

If we have two non-degenerate admissible Hamiltonians $H_1 < H_2$ and two admissible almost complex structures J_1, J_2 , then there is a natural map:

$$SH_*^{(a,b]}(H_1, J_1, \eta) \longrightarrow SH_*^{(a,b]}(H_2, J_2, \eta)$$

This map is called a **continuation map**. This map is defined from a map C on the chain level as follows:

$$C : CF_k^d(H_1, J_1, \eta) \longrightarrow CF_k^d(H_2, J_2, \eta)$$

$$\partial \langle o_- \rangle := \sum_{i(o_-)=i(o_-)} \# P(x_-, x_+) \langle o_+ \rangle$$

where $P(o_-, o_+)$ is a compact oriented zero dimensional manifold of solutions of the following equations: Let K_s be a smooth non-decreasing family of admissible Hamiltonians equal to H_1 for $s \ll 0$ and H_2 for $s \gg 0$ and J_s a smooth family of admissible almost complex structures joining J_1 and J_2 . The set $P(o_-, o_+)$ is the set of solutions to the parameterized Floer equations

$$\partial_s u + J_{s,t}(u(s, t)) \partial_t u = \nabla^{g_t} K_{s,t}$$

such that $u(s, \cdot)$ converges to $x_{\pm}$ as $s \rightarrow \pm\infty$. For a C^∞ generic family (K_s, J_s) this is a compact zero dimensional manifold (if o_-, o_+ have the same relative index with respect to the cylinder C joining them). Again the class $b \in H^2(N, \mathbb{Z}/2\mathbb{Z})$ enables us to orient this manifold. If we have another such non-decreasing family admissible Hamiltonians joining H_1 and H_2 and another smooth family of admissible almost complex structures joining J_1 and J_2 , then the continuation map induced by this second family is the same as the map induced by (K_s, J_s) . The composition of two continuation maps is a continuation map. If we take the direct limit of all these maps with respect to admissible Hamiltonians H ordered by $<$ such that $H|_N < 0$, then we get our symplectic homology groups $SH_*^{(a,b]}(N, \eta)$. We will write $SH_*^\#(N, \eta)$ or $SH_*^\#(H, J, \eta)$ for $SH_*^{[0,\infty)}(N, \eta)$ or $SH_*^{[0,\infty)}(H, J, \eta)$.

Also we will write SH_* instead of $SH_*^{(-\infty, \infty)}$. If we wish to stress which coefficient field we are using, we will write $SH_*^\#(M, \eta, \mathbb{K})$ or $SH_*^\#(H, J, \eta, \mathbb{K})$ if the field is $\mathbb{K}$ for instance. We will also define $SH_*^{\leq b}(M, \eta, \mathbb{K})$ to be the group $SH_*^{(-\infty, b]}(M, \eta, \mathbb{K})$.

We will dealing with other pairs (H, J) that are not necessarily admissible. The definition of symplectic homology $SH_*^{(a,b]}(H, J, \eta)$ is still the same, although (H, J) has to satisfy some conditions to ensure that we have a well defined symplectic homology group. This will be discussed later in Section 4.1. From now on instead of writing $SH_*^{(a,b]}(H, J, \eta)$ we will suppress the term η and just write $SH_*^{(a,b]}(H, J)$ instead when the context is clear. Also

from now on whenever we have a Liouville domain or symplectic manifold then we will assume that we have chosen such a pair $\eta = (\tau, b)$.

2.3. Growth rates. In order to define growth rates, we will need some linear algebra first. Let $(V_x)_{x \in [1, \infty)}$ be a family of vector spaces indexed by $[1, \infty)$. For each $x_1 \leq x_2$ we will assume that there is a homomorphism ϕ_{x_1, x_2} from V_{x_1} to V_{x_2} with the property that for all $x_1 \leq x_2 \leq x_3$, $\phi_{x_2, x_3} \circ \phi_{x_1, x_2} = \phi_{x_1, x_3}$ and $\phi_{x_1, x_1} = \text{id}$. We call such a family of vector spaces a *filtered directed system*. Because these vector spaces form a directed system, we can take the direct limit $V := \varinjlim_x V_x$. From now on we will assume that each V_x is finite dimensional. For each $x \in [1, \infty)$ there is a natural map:

$$q_x : V_x \rightarrow \varinjlim_x V_x.$$

Let $a : [1, \infty) \rightarrow [0, \infty)$ be a function such that $a(x)$ is the rank of the image of the above map q_x . We define the growth rate as:

$$\Gamma((V_x)) := \varlimsup_x \frac{\log a(x)}{\log x} \in \{-\infty\} \cup [0, \infty].$$

If $a(x)$ is 0 then we just define $\log a(x)$ as $-\infty$. If $a(x)$ was some polynomial of degree n with positive leading coefficient, then the growth rate would be equal to n . If $a(x)$ was an exponential function with positive exponent, then the growth rate is ∞ . The good thing about growth rate is that if we had some additional vector spaces $(V'_x)_{x \in [1, \infty)}$ such that the associated function $a'(x) := \text{rank}(V'_x \rightarrow \varinjlim_x V'_x)$ satisfies $a'(x) = Aa(Bx)$ for some constants $A, B > 0$ then $\Gamma(V'_x) = \Gamma(V_x)$. The notation we use for filtered directed systems is usually of the form (V_x) or (V_*) , and we will usually write V_x without brackets if we mean the vector space indexed by x .

In the previous section we defined for a Liouville domain N (whose boundary had discrete period spectrum), $SH_*^{\leq \lambda}(N)$. For $\lambda_1 \leq \lambda_2$, there is a natural map $SH_*^{\leq \lambda_1}(N) \rightarrow SH_*^{\leq \lambda_2}(N)$ given by inclusion of the respective chain complexes. This is a filtered directed system $(SH_*^{\leq \lambda}(N))$ whose direct limit is $SH_*(N)$.

Definition 2.3. We define the growth rate $\Gamma(N, b)$ as:

$$\Gamma(N, b) := \Gamma(SH_*^{\leq \lambda}(N, b))$$

We also have the following Theorem:

Theorem 2.4. Let N_1, N_2 be two Liouville domains such that $\widehat{N}_1$ is symplectomorphic to $\widehat{N}_2$ where the symplectomorphism pulls back $b_2 \in H^2(N_2, \mathbb{Z}/2\mathbb{Z})$ to $b_1 \in H^2(N_1, \mathbb{Z}/2\mathbb{Z})$ and τ_2 to τ_1 where τ_2 and τ_1 are trivializations of the canonical bundle. Then $\Gamma(N_1, (\tau_1, b_1)) = \Gamma(N_2, (\tau_2, b_2))$.

This theorem will be proven in Section 4.1. Hence we will just write $\Gamma(\widehat{N}, d\theta_N, (\tau, b))$ for the growth rate of (N, θ_N) . We will sometimes just write $\Gamma(\widehat{N})$ if the context makes it clear that $d\theta_N$ is our symplectic form and (τ, b) is our associated trivialization and homology class.

3. GROWTH RATE LINEAR ALGEBRA

Recall that a filtered directed system is a family of vector spaces (V_x) parameterized by $[1, \infty)$ forming a category where for $x_1 \leq x_2$ there is a unique homomorphism from V_{x_1} to V_{x_2} and no other morphisms anywhere else. For technical reasons we will define V_x to be zero for $x < 1$ and so all the morphisms starting with one of these vector spaces is also 0. A morphism of filtered directed systems $\phi : (V_x) \rightarrow (V'_x)$ consists of some constant C_ϕ and a sequence of maps

$$a_x : V_x \rightarrow V'_{C_\phi x}$$

so that we have the following commutative diagram:

$$\begin{array}{ccc} V_{x_1} & \xrightarrow{a_{x_1}} & V'_{C_\phi x_1} \\ \downarrow & & \downarrow \\ V_{x_2} & \xrightarrow{a_{x_2}} & V'_{C_\phi x_2} \\ \downarrow & & \downarrow \\ V_{x_3} & \xrightarrow{a_{x_3}} & V'_{C_\phi x_3} \end{array}$$

for all $x_1 \leq x_2 \leq x_3$ where the vertical arrows come from the directed system.

Let ψ_{x_1, x_2} be the natural map from V_{x_1} to V_{x_2} in this filtered directed system for $x_1 \leq x_2$. For each constant $C \geq 0$, we have an morphism C_V from (V_x) to (V_x) given by the map $\psi_{x, Cx}$. We say that (V_x) and (V'_x) are *isomorphic* if there is a morphism ϕ from (V_x) to (V'_x) and another morphism ϕ' from (V'_x) to (V_x) such that $\phi' \circ \phi = C_V$ and $\phi \circ \phi' = C'_{V'}$, where $C, C' \geq 0$ are constants and $C_V : (V_x) \rightarrow (V_x)$, $C'_{V'} : (V'_x) \rightarrow (V'_x)$ are the morphisms described above. One of the aims of this paper is to assign for each completion $\widehat{M}$ of a Liouville domain M , a filtered directed system which unique up to isomorphism. From this we can define growth rate. In order for growth rate to be well defined, we need to show:

Lemma 3.1. *Let $(V_x), (V'_x)$ be two isomorphic filtered directed systems, then $\Gamma(V_x) = \Gamma(V'_x)$.*

Proof. Let $\phi_x : V_x \rightarrow V'_{C_\phi x}$ and $\phi'_x : V'_x \rightarrow V_{C_{\phi'} x}$ be our isomorphisms. We have a morphism from $\varinjlim_x V_x$ to $\varinjlim_x V'_x$ induced from ϕ and an inverse induced from ϕ' . This is because the morphism induced by C_V is the identity map on $\varinjlim_x V_x$ and similarly $C'_{V'}$ induces the identity map. We will write ϕ and ϕ' for such maps by abuse of notation. Let $a_x : V_x \rightarrow \varinjlim_x V_x$ and $a'_x : V'_x \rightarrow \varinjlim_x V'_x$. Because ϕ is an isomorphism on $\varinjlim_x V_x$, we have that the rank of the image of $\phi \circ a_x$ is equal to the rank of the image of a_x . We have

that $a'_{C_\phi x} \circ \phi_x = \phi \circ a_x$ which implies that $\text{rank im}(a'_{C_\phi x}) \geq \text{rank im}(a_x)$. Similarly $\text{rank im}(a_{C_{\phi'} x}) \geq \text{rank im}(a'_x)$ for all $x \in [1, \infty)$. Hence:

$$\begin{aligned} \varlimsup_x \frac{\log \text{rank im}(a_x)}{\log x} &\leq \varlimsup_x \frac{\log \text{rank im}(a'_{C_\phi x})}{\log x} = \varlimsup_x \frac{\log \text{rank im}(a'_{C_\phi x})}{\log C_\phi x} = \\ \varlimsup_x \frac{\log \text{rank im}(a'_x)}{\log x} &\leq \varlimsup_x \frac{\log \text{rank im}(a_{C_{\phi'} x})}{\log x} = \varlimsup_x \frac{\log \text{rank im}(a_x)}{\log x}. \end{aligned}$$

This implies that $\Gamma(V_x) = \Gamma(V'_x)$ as the first term in the above set of inequalities is $\Gamma(V_x)$ and the fourth term is $\Gamma(V'_x)$. $\square$

We now need a Lemma giving us a sufficient condition telling us when two filtered directed systems are equivalent.

Lemma 3.2. *Let (V_x^j) ($j = 1, 2, 3, 4$) be filtered directed systems. Let $u_j : V_x^j \rightarrow V_{C_j x}^{j+1}$ ($j = 1, 2, 3$) be morphisms of directed systems so that composing any two of them gives us an isomorphism. Then V_x^2 is isomorphic to V_x^3 .*

Proof. of Lemma 3.2. Let ψ_{x_1, x_2}^j be the directed system map $V_{x_1}^j \rightarrow V_{x_2}^j$ for $x_1 \leq x_2$. By the definition of an isomorphism of filtered directed systems, there exist morphisms $b_1 : V_x^3 \rightarrow V_{D_1 x}^1$ and $b_2 : V_x^4 \rightarrow V_{D_2 x}^2$ so that $u_2 \circ u_1 \circ b_1$ is the directed system map

$$\psi_{x, D_1 C_1 C_2 x}^3 : V_x^3 \rightarrow V_{D_1 C_1 C_2 x}^3.$$

Also

$$b_1 \circ u_2 \circ u_1 = \psi_{x, D_1 C_1 C_2 x}^1,$$

$$u_3 \circ u_2 \circ b_2 = \psi_{x, D_2 C_2 C_3 x}^4$$

and

$$b_2 \circ u_3 \circ u_2 = \psi_{x, D_2 C_2 C_3 x}^2.$$

We define $\phi : V_x^2 \rightarrow V_{C_2 x}^3$ by $\phi = u_2$. We define $\phi' : V_x^3 \rightarrow V_{D_1 D_2 C_1 C_2 C_3 x}^2$ by $\phi' = b_2 \circ u_3 \circ \psi_{x, D_1 C_1 C_2 x}^3$.

Let $x \in V_x^3$. By abuse of notation for any x we will just write ψ_K^j for $\psi_{x, Kx}^j$ for $j = 1, 2, 3$. Because $\psi_{D_1 C_1 C_2}^3 = u_2 \circ u_1 \circ b_1$,

$$\begin{aligned} \phi \circ \phi' &= u_2 \circ b_2 \circ u_3 \circ u_2 \circ u_1 \circ b_1 \\ &= u_2 \circ \psi_{D_2 C_2 C_3}^2 \circ u_1 \circ b_1 = \psi_{D_2 C_2 C_3}^3 \circ u_2 \circ u_1 \circ b_1 \\ &= \psi_{x, D_2 C_2 C_3 x}^3 \circ \psi_{D_1 C_1 C_2}^3. \end{aligned}$$

Also:

$$\begin{aligned} \phi' \circ \phi &= b_2 \circ u_3 \circ \psi_{D_1 C_1 C_2}^3 \circ u_2 \\ &= \psi_{D_1 C_1 C_2}^2 \circ b_2 \circ u_3 \circ u_2 \\ &= \psi_{D_1 C_1 C_2}^2 \circ \psi_{D_2 C_2 C_3}^2. \end{aligned}$$

Hence ϕ and ϕ' give us our isomorphism and we have proven the Lemma. $\square$

4. GROWTH RATE GEOMETRY

4.1. Some alternate definitions of growth rate. We will define growth rate for a slightly larger class of manifolds called finite type convex symplectic manifolds and also using a broader class of Hamiltonians. There are three reasons for doing this. The first reason is that we wish to prove that growth rate is an invariant up to symplectomorphism and so we need a definition of growth rate which is invariant under symplectomorphism. The second reason is that the author wishes to use this larger class of Hamiltonians in a future paper to prove that growth rate behaves well under products and also with respect to Lefschetz fibrations. A third reason is that this way of thinking might be useful for answering various dynamical questions.

A *convex symplectic manifold* is a manifold M with a 1-form θ_M such that

- (1) $\omega_M := d\theta_M$ is a symplectic form.
- (2) There is an exhausting function $f_M : M \rightarrow \mathbb{R}$ and a sequence $c_1 < c_2 < \dots$ tending to infinity such that the ω_M -dual X_{θ_M} of θ_M satisfies $df_M(X_{\theta_M}) > 0$ along $f_M^{-1}(c_i)$ for each i .

Some basic facts about convex symplectic manifolds are proven in the appendix. We say that M is of *finite type* if there is a $C \in \mathbb{R}$ such that $(f_M^{-1}(-\infty, c], \theta_M)$ is a Liouville domain for all $c \geq C$.

Let (M, θ_M^t) be a smooth family of convex symplectic manifolds parameterized by $t \in [0, 1]$. This is said to be a *convex deformation* if for every $t \in [0, 1]$ there is a $\delta_t > 0$ and an exhausting function f_M^t and a sequence of constants $c_1^t < c_2^t < \dots$ tending to infinity such that $((f_M^t)^{-1}(-\infty, c_i^t], \theta_M^t)$ is a Liouville domain for each $s \in [t - \delta_t, t + \delta_t]$ and each i . We do not require that f_M^t, c_i^t, δ_t smoothly varies with t . In fact it can vary in a discontinuous way with t .

Let M be a finite type convex symplectic manifold. In order to define growth rate, we need a slightly larger class of Hamiltonians. We will first describe Hamiltonians on M that give us filtered directed systems. Let (S, j) be a complex surface possibly with boundary. Let γ be a 1-form on S so that $d\gamma \geq 0$. Let $H : M \rightarrow \mathbb{R}$ be a Hamiltonian and J an almost complex structure compatible with the symplectic form ω . We have a Hamiltonian vector field X_H defined by $\omega(X_H, \cdot) = dH$. A map $u : S \rightarrow M$ satisfies the *perturbed Cauchy-Riemann equations* if $(du - X_H \otimes \gamma)^{0,1} = 0$. Here $du - X_H \otimes \gamma$ is a 1-form on S with values in the complex vector bundle $\text{Hom}(TS, u^*TM)$ where the complex structure at a point $s \in S$ is induced from j and J . The equation $(du - X_H \otimes \gamma)^{0,1} = 0$ is written explicitly as

$$(1) \quad du - X_H \otimes \gamma + J \circ (du - X_H \otimes \gamma) \circ j = 0.$$

Sometimes we will write $(du - X_H \otimes \gamma)_J^{0,1} = 0$ if we wish to emphasise the fact that we are using the almost complex structure J . Here is a particular example. Let $S = \mathbb{R} \times S^1 = \mathbb{C}/\mathbb{Z}$. We let $\gamma = dt$ where t parameterizes $S^1 = \mathbb{R}/\mathbb{Z}$. Then the perturbed Cauchy-Riemann equations become

$$\partial_s u + J \partial_t u = J X_H$$

which is just the Floer equation.

The pair (H, J) on M is said to satisfy a *maximum principle with respect to an open set U^H* if there is a compact set $K' \subset M$ containing U^H such that for any compact complex surface (S, j) with 1-form γ ($d\gamma \geq 0$) and map $u : S \rightarrow M$, satisfying:

- (1) u satisfies the perturbed Cauchy-Riemann equations.
- (2) $u(\partial S) \subset U^H$

we have that $u(S) \subset K'$. We also require that U^H contains all the 1-periodic orbits of (H, J) of action greater than some small negative constant.

A pair (H, J) is said to be SH_* admissible if there is a discrete subset $A_H \subset (0, \infty)$ such that $(\lambda H, J)$ satisfies the maximum principle for $\lambda \in (0, \infty) \setminus A_H$ with respect to some relatively compact open set U_λ^H . We require that $U_{\lambda_1}^H \subset U_{\lambda_2}^H$ for $\lambda_1 \leq \lambda_2$. Note that if (H, J) is SH_* admissible then so is any positive multiple of H .

For an SH_* admissible pair (H, J) , we define $SH_*^\#(\lambda H, J)$ ($\lambda \in (0, \infty) \setminus A_H$) as follows: By Lemma 2.2, we can perturb λH by an arbitrarily small amount to a non-degenerate time dependent Hamiltonian $H'_t : S^1 \times M \rightarrow \mathbb{R}$ so that it is equal to λH outside some closed subset of U_λ^H . After subtracting a small constant from H'_t , we can assume that H'_t is equal to $\lambda H - \epsilon$ on a closed subset of U_λ^H and $H'_t < \lambda H$ where $\epsilon > 0$ is some constant. For a generic S^1 family J'_t of almost complex structures such that $J_t = J$ outside a closed subset of U_λ^H , we have that $SH_*^\#(H'_t, J'_t)$ is well defined. This is because all the 1-periodic orbits of H'_t are contained in a compact set and the maximum principle ensures that all the Floer trajectories also stay inside a compact set. The pair (H'_t, J'_t) constructed above is called an *approximating pair* for (H, J) . A similar argument ensures that if we have two approximating pairs (H'_t, J'_t) and (H''_t, J''_t) for (H, J) with $H'_t < H''_t$, there is a well defined continuation map $SH_*^\#(H'_t, J'_t) \rightarrow SH_*^\#(H''_t, J''_t)$ induced from a generic family of approximating pairs joining (H'_t, J'_t) and (H''_t, J''_t) where the Hamiltonians are non-decreasing. Also because the continuation map is induced from a non-decreasing family of Hamiltonians, we have that the continuation map respects the filtration by action. So any orbit of action $\leq f$ is sent to another orbit of action $\leq f$ under the continuation map. The set of approximating pairs (H'_t, J'_t) induces a directed system where $(H'_t, J'_t) \leq (H''_t, J''_t)$ if and only if $H'_t \leq H''_t$ for all $t \in S^1$. We define $SH_*^\#(\lambda H, J)$ as $\varinjlim_{(H'_t, J'_t)} SH_*^\#(H'_t, J'_t)$ where the direct limit is taken over this directed system.

We also have a continuation map for $\lambda_1 \leq \lambda_2$ where $\lambda_i \in (0, \infty) \setminus A_H$ from $SH_*^\#(\lambda_1 H, J)$ to $SH_*^\#(\lambda_2 H, J)$. This continuation map is induced by a family of pairs (H_t^s, J_t^s) such that

- (1) they are equal to $(\lambda'_s H - \epsilon_s, J)$ outside a closed subset of $U_{\lambda_2}^H$ for some smooth non-decreasing family of constants λ'_s and some smooth non-increasing family of constants ϵ_s .

- (2) (H_t^s, J_t^s) is a fixed approximating pair (H_t^-, J_t^-) for $(\lambda_1 H, J)$ for $s \ll 0$.
- (3) (H_t^s, J_t^s) is a fixed approximating pair (H_t^+, J_t^+, t) for $(\lambda_2 H, J)$ for $s \gg 0$.
- (4) $\frac{\partial}{\partial s} H_t^s \geq 0$.

We need to show that the continuation map trajectories stay inside a compact set. Suppose we have some map $u : \mathbb{R} \times S^1 \rightarrow M$ satisfying the continuation map equations which join orbits inside $U_{\lambda_2}^H$:

$$\partial_s u + J_t^s \partial_t u = J_t^s X_{H_t^s}.$$

These can be rewritten in the following way:

$$(du - X_{H_t^s} \otimes dt)_{J_t^s}^{0,1} = 0.$$

We have $(H_t^s, J_t^s) = (\lambda'_s H - \epsilon_s, J)$ outside some compact subset R of $U_{\lambda_2}^H$. Let $S \subset \mathbb{R} \times S^1$ be a compact submanifold so that $u(S)$ is disjoint from R and $u(\partial S) \subset U_{\lambda_2}^H$. Hence we have that u restricted to S satisfies:

$$(du - X_{\lambda'_s H - \epsilon_s} \otimes dt)_J^{0,1} = 0.$$

The constants ϵ_s do not matter so our equation becomes:

$$(du - X_H \otimes \gamma)_J^{0,1} = 0$$

where $\gamma = \lambda'_s dt$. Here $d\gamma \geq 0$ because λ'_s is non-decreasing. Hence by the maximum principle we have that S must be contained entirely inside $U_{\lambda_2}^H$. This implies that the image of each continuation map u must be contained inside $U_{\lambda_2}^H$ and hence inside a fixed compact set. Hence we have a well defined continuation map between $SH_*^\#(H_t^-, J_t^-)$ and $SH_*^\#(H_t^+, J_t^+)$. This induces a continuation map between $SH_*^\#(\lambda_1 H, J)$ and $SH_*^\#(\lambda_2 H, J)$. If $\lambda \in A_H$ then we define $SH_*^\#(\lambda H, J)$ as the direct limit of $SH_*^\#(\lambda' H, J)$ where $\lambda' \notin A$ tends to λ from below. Hence we have a filtered directed system $(SH_*^\#(\lambda H_t, J_t))$.

We wish to put some additional conditions on the pair (H, J) so that the associated filtered directed system is an invariant related to symplectic homology. Here are the additional conditions:

- (1) (bounded below property)

The Hamiltonian H is greater than or equal to zero, and there exists a compact set K and a constant $\delta_H > 0$ such that: $H > \delta_H$ outside K .

- (2) (Liouville vector field property)

There exists an exhausting function f_H , and 1-form θ_H such that:

- (a) $\theta_M - \theta_H$ is exact where θ_M is the Liouville form on M .
- (b) There exists a small $\epsilon_H > 0$ such that $dH(X_{\theta_H}) > 0$ in the region $H^{-1}(0, \epsilon_H]$ where X_{θ_H} is the ω_M -dual of θ_H . We also require that $d(dH(X_{\theta_H}))(X_{\theta_H}) > dH(X_{\theta_H})$ in this region.

- (c) There is a constant C such that $df_H(X_{\theta_H}) > 0$ in the region $f_H^{-1}[C, \infty)$ and $f_H^{-1}(-\infty, C]$ is non-empty and is contained in the interior of $H^{-1}(0)$.
- (3) (action bound property)

There is a constant C_H and 1-form θ such that the function $-\theta(X_H) - H$ must be bounded above by C_H where X_H is the Hamiltonian vector field associated to H_t . We also require that $\theta - \theta_M$ is exact.

If an SH_* admissible pair (H, J) has these additional conditions, then it is called *growth rate admissible*.

Here are some important examples:

First example : Let X_{θ_M} be the vector field given by the ω_M -dual of θ_M . Because M is a finite type convex symplectic manifold, we have a function $f_M : M \rightarrow \mathbb{R}$ such that, $df_M(X_{\theta_M}) > 0$ in the region where $f_M > c$ for some $c \gg 0$ and such that f_M is exhausting. Let X_{f_M} be the Hamiltonian vector field of f_M . Choose some $c < c_1 < c_2 < \dots$ tending to infinity so that $df_M(X_{\theta_M}) > 0$ on $f_M^{-1}(c_i)$ for each i . We perturb f_M by a C^0 small amount to g_M so that it has the following property: $g_M^{-1}(c_i) = f_M^{-1}(c_i)$ and $dg_M(X_{\theta_M})$ is constant on a small neighbourhood of $g_M^{-1}(c_i)$ for each i and $dg_M(X_{\theta_M}) > 0$ for $g_M > c$. The level sets $A_y := g_M^{-1}(y)$ for $y > c$ are contact manifolds with contact form $\alpha_y := \theta_M|_{A_y}$. Because

$$\alpha_y(X_{g_M}) = \theta_M(X_{g_M}) = \omega_M(X_{\theta_M}, X_{g_M}) = -dg_M(X_{\theta_M}) < 0,$$

we have that X_{g_M} is non-trivial in the region $\{g_M > c\}$. Because the vector field X_{g_M} is contained in TA_y ($y > c$), and because it is non-zero, it has a shortest orbit inside A_y (as A_y is compact). The reason for this is that for each $p \in A_y$ we can choose coordinates $x_1, \dots, x_{2n}$ around p so that $X_{g_M} = \frac{\partial}{\partial x_1}$, which means that any flowline going through a smaller neighbourhood U of p must take some time $\delta_U > 0$ to pass through U . Because A_y is compact, we can then cover it with finitely many such neighborhoods U_i which implies that any orbit must flow for time at least $\min_i \delta_{U_i}$. Let $\delta(y) > 0$ be smaller than $\min_i \delta_{U_i}$. We can assume that $\delta(y)$ is a smooth function of y .

Let $h : \mathbb{R} \rightarrow \mathbb{R}$ be a function with the property that $h(x) = 0$ for $x \leq c+1$ and $h(x) > 0$ elsewhere. For $x \geq c+1$ we let $h'(x)$ be smaller than $\delta(x)/g_M$ and $h' > 0$ whenever $h > 0$. We also assume that h' is small enough so that $h'(x)\alpha(x)(X_{g_M})$ is bounded above by a constant. The Hamiltonian flow of $H := h(g_M)$ in the region $g_M \leq c$ is 0 and for $x > c+1$ in the region A_x , it is equal to some very small multiple of $X_{g_M}|_{A_x}$. We also need an additional technical condition which ensures that our Hamiltonian H will satisfy the Liouville vector field property. We require that there exists some $\epsilon_h > 0$ with $h''(x) > 0$ for $x \in (c+1, c+1+\epsilon_h)$. Fix $\lambda > c$. In the region $\{g_M > \lambda\}$, there are no 1-periodic orbits of λH . This is because all the orbits of $X_H|_{A_x} = h'(x)X_{g_M}|_{A_x}$ have length greater than λ because

$h' < \delta_x/g_M \leq \delta_x/\lambda$. Also because $H \geq 0$ and $\alpha_x(X_H)$ is bounded from above for $x > c$, we have that $-\theta_M(X_H) - H$ is bounded from above so H satisfies the action bound property. The Hamiltonian H satisfies the bounded below property because its zero set is compact, it is greater than or equal to zero and is greater than some constant near infinity. Note that $\log(h'(x))$ tends to minus infinity as x tends to $c + 1$ from above. Also the derivative of this function is positive. This implies that the derivative $h''(x)/h'(x)$ tends to infinity. Hence (after shrinking ϵ_h) we can assume that $h''(x) \geq \nu h'(x)$ for $x \in (c + 1, c + 1 + \epsilon_h)$ where


$$\nu > (1 - d(g_M(X_{\theta_M}))(X_{\theta_M})) / g_M(X_{\theta_M})^2$$

in the region $x \in (c + 1, c + 1 + \epsilon_h)$. Because $dH(X_{\theta_M}) = h'(g_M)dg_M(X_{\theta_M}) \geq 0$ and

$$\begin{aligned} d(dH(X_{\theta_M}))(X_{\theta_M}) &= h''(g_M)dg_M(X_{\theta_M})^2 + h'(g_M)d(g_M(X_{\theta_M}))(X_{\theta_M}) > \\ &h'(g_M)dg_M(X_{\theta_M}) \end{aligned}$$

in the region $H^{-1}(0, \epsilon_H)$ for some small $\epsilon_H > 0$, H satisfies the Liouville vector field property. We need to find a J so that (H, J) satisfies a maximum principle. Because $dg_M(X_{\theta_M})$ is constant on a small neighbourhood of c_i for each i , we have (by flowing G_i along X_{θ_M}) a small neighbourhood of $G_i := g_M^{-1}(c_i)$ symplectomorphic to $(1 - \epsilon_i, 1 + \epsilon_i) \times G_i$ with contact form $\kappa_i g_M \alpha_i$ where $\alpha_i := \theta_M|_{G_i}$ is a contact form and κ_i is a constant. This is a slice of a positive cylindrical end so we can choose J so that it looks cylindrical in these regions. We define U_λ^H to be any family open sets of the form $g_M^{-1}(-\infty, c_i)$ containing all the 1-periodic orbits of λH and such that $U_{\lambda_1}^H \subset U_{\lambda_2}^H$ for $\lambda_1 \leq \lambda_2$. Hence by Lemma 9.1, there exists a J such that $(\lambda H, J)$ satisfies the maximum principle with respect to U_λ^H . Hence (H, J) is growth rate admissible.

Second example : Let f_M and X_{θ_M} be as above. Let $Q := f_M^{-1}(-\infty, C]$ be a manifold with boundary such that X_{θ_M} is transverse to the boundary of Q and pointing outwards and $df_M(X_{\theta_M}) > 0$ outside Q . Even though the Liouville vector field X_{θ_M} may not be complete, we can still flow ∂Q along X_{θ_M} so that we have that the set $\overline{(M \setminus Q)}$ is diffeomorphic to some open subset U of $[1, \infty) \times \partial Q$ containing $\{1\} \times \partial Q$. We have that $\theta_M|_U = r\alpha$ where r parameterizes $[1, \infty)$ and $\alpha = \theta_M|_{\partial Q}$. We call this a *partial cylindrical end* of M . We can ensure that the period spectrum of the contact manifold ∂Q is discrete and injective (after perturbing f_M very slightly). Let J be an almost complex structure which is cylindrical inside U (i.e. $dr \circ J = -\theta_M$). Let $\epsilon > 0$ be a constant small enough so that $r^{-1}(1 + \epsilon)$ is still a compact manifold transverse to X_{θ_M} . Let H be a Hamiltonian such that $H = h(r)$ in U and 0 inside Q . We require that $h(r) = 0$ near $r = 1$ and $h(r) = r - \frac{\epsilon}{4}$ for $r \geq 1 + \epsilon/2$, and $h', h'' \geq 0$ and $h' > 0$ when $h > 0$. We also require that $h''(x) > 0$ for $x \in (1, 1 + \epsilon_H)$ where $\epsilon_H > 0$ is small.



For λ not in the period spectrum of ∂Q , we have that all of the 1-periodic orbits of λH lie inside the compact set $\{r \leq \epsilon/2\}$. We define U_λ^H to be the open set $Q \cup r^{-1}([1, 1 + \epsilon])$ for all λ . The pair $(\lambda H, J)$ satisfies the maximum principle with respect to U_λ^H by 9.1. Hence (H, J) is SH_* admissible. We have that (H, J) satisfies the bounded below property and the Liouville vector field property because $X_{\theta_M} = r \frac{\partial}{\partial r}$ in U , $h' > 0$ when $h > 0$ and $h''(r) > 0$ for $r \in (1, 1 + \epsilon_H)$. We also have that (H, J) satisfies the action bound property because $-\theta_M(X_H) - H = rh'(r) - r + \frac{\epsilon}{4} = \frac{\epsilon}{4}$ outside a compact set (where $H = r - \frac{\epsilon}{4}$). All of this means that (H, J) is growth rate admissible.

Suppose that $(H_0, J_0), (H_1, J_1)$ is growth rate admissible with the property that $U_\lambda^{H_0} \subset U_\lambda^{H_1}$ for all λ and such that $(H_1, J_1) = (\kappa H_0 + c_1, J_0)$ outside a closed subset of $U_\lambda^{H_0}$. Suppose we have a non-decreasing family of pairs (H_t, J_t) equal to $(\kappa_t H_0 + c_t, J)$ outside a closed subset of $U_\lambda^{H_0}$ for some smooth family of constants $\kappa_t \geq 1, c_t \in \mathbb{R}$. Then we have a well defined continuation map from $SH_*^\#(\lambda H_0, J_0)$ to $SH_*^\#(\lambda H_1, J_1)$ and this induces a morphism of filtered directed systems. We call such a morphism a *restricted continuation morphism*. Suppose now we have two growth rate admissible pairs (H'_0, J'_0) and (H'_1, J'_1) with the following properties:

- (1) $U_\lambda^{H'_0} = U_\lambda^{H'_1}$.
- (2) $(H'_0, J'_0) = (H'_1, J'_1)$ on a neighbourhood of $\cup_\lambda U_\lambda^{H'_0}$.

Then the filtered directed systems $(SH_*^\#(\lambda H'_0, J'_0))$ and $(SH_*^\#(\lambda H'_1, J'_1))$ are isomorphic because all 1-periodic orbits of non-negative action and all Floer trajectories connecting them are identical. We call such a morphism a *switch isomorphism*.

The main theorem of this section is:

Theorem 4.1. *Let $(H, J), (H', J')$ be growth rate admissible. Then the filtered directed system $(SH_*^\#(\lambda H, J))$ is isomorphic to $(SH_*^\#(\lambda H', J'))$ as filtered directed systems.*

The isomorphism between these filtered directed systems is a composition of restricted continuation morphisms and switch morphisms and inverses of these morphisms.

The reason why we mention restricted continuation morphisms and switch morphisms is because we would like this result to hold if $SH_*^\#$ had some additional algebraic structure as well such as the pair of pants product so that it can be used in future work. The point is that if restricted continuation morphisms and switch morphisms are also morphisms preserving this additional algebraic structure then we immediately get an invariance result for symplectic homology with this extra structure.

We need some preliminary lemmas before we prove this Theorem.

Lemma 4.2. *Let H be a Hamiltonian satisfying the Liouville vector field property, then there is a growth rate admissible pair (H^p, J^p) such that:*

- (1) $H^p = H$ on a small neighbourhood of $H^{-1}(0)$ and $(H^p)^{-1}(0) = H^{-1}(0)$.
- (2) $-\theta_H(X_{H^p}) - H^p \geq 0$ everywhere and $-\theta_H(X_{H^p}) - H^p > 0$ when $H^p > 0$.
- (3) $-\theta_H(X_{H^p}) - H^p$ is greater than some constant $\delta'_{H^p} > 0$ outside a large compact set.
- (4) The construction of H^p only depends on H near $H^{-1}(0)$.
- (5) There is a fixed compact set K' such that all 1-periodic orbits of λH^p are contained in the interior of K' . This set is an embedded codimension 0 manifold with boundary and we can ensure that it fits inside any open set containing $H^{-1}(0)$.
- (6) Any solution $u : S \rightarrow M$ of the perturbed Cauchy-Riemann equations with respect to $(\tilde{H}, \tilde{J})$ where $\tilde{H} > 0$, $(\tilde{H}, \tilde{J}) = (\lambda H^p, J)$ inside K' and $u(\partial S) \subset K'$ is contained in K' .

Here θ_H is the 1-form that makes H satisfy the Liouville vector field property.

Note that this Lemma also tells us the following fact about H : we have that $-\theta_H(H) - H > 0$ on some small neighbourhood of $H^{-1}(0)$. This will be useful later on.

Proof. of Lemma 4.2. The main idea of this proof is to use bump functions to extend (H, J) restricted to a small neighbourhood of $H^{-1}(0)$ to a pair which looks like the pair described in the second example. Here θ_H is a 1-form such that $\theta_H - \theta_M$ is an exact 1-form. The vector field X_{θ_H} which is the ω_M -dual of θ_H has the property that there is a function $\underline{f_H}$ such that $d\underline{f_H}(X_{\theta_H}) > 0$ inside a small neighbourhood of the $\overline{M} \setminus H^{-1}(0)$. We also have an ϵ_H such that $X_{\theta_H}(H) > 0$ on $H^{-1}(0, \epsilon_H)$ by the Liouville vector field property.

Because we are only interested in what H is near $H^{-1}(0)$ it can be anything we like outside a neighbourhood of this set. So from now on (after changing H outside a neighbourhood of $H^{-1}(0)$) we can assume that H satisfies the bounded below property. Let $Q := f_H^{-1}(-\infty, C]$ be a manifold with

boundary such that X_{θ_H} is transverse to the boundary of Q and pointing outwards and $df_H(X_{\theta_H}) > 0$ outside Q and such that $H^{-1}(0)$ contains Q . We have that ∂Q is a contact manifold with contact form $\alpha = \theta_H|_{\partial Q}$. We flow ∂Q along X_{θ_H} so that we have that the set $\overline{(M \setminus Q)}$ is diffeomorphic to some open subset U of $[1, \infty) \times \partial Q$. This is our partial cylindrical end. We have that $\theta_H|_U = r\alpha$ where r parameterizes $[1, \infty)$, and $X_{\theta_H} = r\frac{\partial}{\partial r}$ in the region U . We have:

$$-\theta_H(X_H) - H = r\frac{\partial H}{\partial r} - H$$

inside U . Because H satisfies the Liouville vector field property we have on a neighbourhood of $H^{-1}(0)$, $d(dH(X_H))(X_H) > dH(X_H) > 0$ outside $H^{-1}(0)$. Because $X_H = r\frac{\partial}{\partial r}$ on our partial cylindrical end this condition becomes:

$$r\frac{\partial H}{\partial r} + r^2\frac{\partial^2 H}{\partial r^2} > r\frac{\partial H}{\partial r} > 0$$

which in turn is equivalent to $\frac{\partial H}{\partial r}, \frac{\partial^2 H}{\partial r^2} > 0$. We shrink ϵ_H so that $H^{-1}[0, \epsilon_H]$ is contained in V . We also have that

$$r\frac{\partial^2 H}{\partial r^2} = \frac{\partial}{\partial r}(-\theta_H(X_H) - H).$$

Hence we have that $-\theta_H(X_H) - H > 0$ in the region $H^{-1}(0, \epsilon_H)$. By the bounded below property, there exists a constant $\delta_H > 0$ such that $H > \delta_H$ outside V . We shrink ϵ_H so that it is smaller than δ_H . Choose a function $q_1 : \mathbb{R} \rightarrow \mathbb{R}$ such that $q_1(x) = 1$ for $x \leq 2\epsilon_H/3$ and $q_1(x) = 0$ for $x \geq \epsilon_H$. Because $dH(X_{\theta_H}) > 0$ on the level set $W := H^{-1}(\epsilon_H/2)$, we have that this level set is regular and is a contact manifold with contact form $\alpha_2 := \theta_H|_W$. There exists a function $f : \partial Q \rightarrow [1, \infty)$ such that under the identification of $\overline{M \setminus Q}$ with a subset of $[1, \infty) \times \partial Q$,

$$W = \{(f(x), x) | x \in \partial Q\} \subset [1, \infty) \times \partial Q.$$

We will write this set as $(r/f)^{-1}(1)$. We also have a new partial cylindrical end which is the region $\{(r/f) \geq 1\}$. This is diffeomorphic to some codimension 0 submanifold with boundary of $W \times [1, \infty)$ where the cylindrical coordinate is (r/f) and the contact form is $\theta_H|_W$. Let q_2 be a function on $\mathbb{R}$ satisfying:

- (1) $q_2(x) = 0$ for $x \leq 0$
- (2) $q_2'(x), q_2''(x) \geq 0$.
- (3) We let $q_2'(x)$ be constant and equal to 1 for $x \geq \delta_r$. Here δ_r is a constant such that the level sets $(r/f)^{-1}(x)$ are all compact for $1 \leq x \leq \delta_r + 1$. So for $x \geq \delta_r$, $q_2 = r - \iota_r$ where $\iota_r > 0$ is a constant.
- (4) For $x > 0$ we also choose q_2 so that $q_2'(x) > 0$.

Let $H^p := q_1(H)H + \kappa q_2(r/f - 1)$ where $\kappa > 0$ is a constant to be determined. Here $q_1(H)$ has support inside the region V because $H \geq \epsilon_H$ outside V . We have for $\kappa \gg 0$, that $-\theta_H(H^p)(X_{\theta_H}) - H^p \geq 0$. This is because $H^p =$

$H + \kappa q_2(r/f - 1)$ in $H^{-1}(0, 2\epsilon_H/3)$ and $-\theta_H(H) - H > 0$, $-\theta_H(q_2(r/f - 1)) - q_2(r/f - 1) \geq 0$ in this region because $q_2, q_2', q_2'' \geq 0$. Also $-\theta_H(r/f - 1 - \iota_r) - r/f + \iota_r = \iota_r$ is greater than some fixed constant outside $H^{-1}(-\infty, 2\epsilon_H/3)$ hence by the action bound property we get that $-\theta_H(X_{H^p}) - H^p \geq 0$ for κ large enough. Hence in the region where $H^p > 0$, we have that $-\theta_H(X_{H^p}) - H^p > 0$.

We also let J^p be an almost complex structure such that it is cylindrical on the partial cylindrical end $\{(r/f) \geq 1\}$ viewed as a subset of $W \times [1, \infty)$. We can perturb f very slightly so that the period spectrum of $W = (r/f)^{-1}(1)$ is discrete. Define $K' := M \setminus (r/f)^{-1}(1, \infty)$. This is a compact co-dimension 0 manifold whose boundary is W . Because the period spectrum of W is discrete and equal to $A \subset (0, \infty)$, we have for all $\lambda \in (0, \infty) \setminus A$, λH^p has all its 1-periodic orbits contained in a compact subset of M . Also 9.1 with boundary inside K' must stay in K' . This implies that (H^p, J^p) satisfies the maximum principle. We also have that because $H^p \geq 0$ and $H^p = (r/f) - \iota_r$ near infinity, H^p satisfies the bounded below property. Because H satisfies the Liouville vector field property, then so does H^p as $H^p = H$ on a small neighbourhood of $H^{-1}(0)$ and $H^{p-1}(0) = H^{-1}(0)$. We also have that it satisfies the action bound property because $-\theta_H(r/f - \iota_r) - (r/f - \iota_r) = \iota_r$ near infinity.

Finally property (6) is satisfied by Lemma 9.1 as well. $\square$

Before we prove Theorem 4.1 we will prove a couple of weaker versions of it. Here is one.

Lemma 4.3. *Let $(H, J), (H', J')$ be growth rate admissible pairs satisfying the following properties:*

- (1) $H' = H$ in a neighbourhood of $H^{-1}(0)$.
- (2) $H^{-1}(0) = H'^{-1}(0)$.
- (3) $U_\lambda^H = U_\lambda^{H'}$.
- (4) $(H, J) = (H' + c, J')$ outside a closed subset of $U_1^H = \cap_\lambda U_\lambda^H$.

Then there is a restricted continuation morphism from $(SH_*^\#(\lambda H, J))$ to $(SH_*^\#(\lambda H', J'))$ which is an isomorphism. Its inverse is also a restricted continuation morphism.

Proof. of Lemma 4.3. By the bounded below property there exists a constant $\delta > 0$ with $H, H' > \delta$ outside a compact set K . This implies that there exists constants $Q_1, Q_2 > 1$ such that $H \leq Q_1 H'$ and $H' \leq Q_2 H$. Let (H^s, J^s) ($s \in \mathbb{R}$) be a family of pairs such that

- (1) $(H^s, J^s) = (H, J)$ for $s \ll 0$ and $(H^s, J^s) = (Q_1 H', J')$ for $s \gg 0$.
- (2) We let $(H^s, J^s) = (\kappa_s H + c_s, J)$ outside a closed subset of $\cap U_\lambda^H$ for some non-decreasing family of constants $\kappa_s \geq 1$, $c_s \in \mathbb{R}$.
- (3) $\frac{\partial H^s}{\partial s} \geq 0$.

The family of pairs $(\lambda H^s, J^s)$ induces a continuation map ϕ from the group $SH_*^\#(\lambda H, J)$ to $SH_*^\#(\lambda Q_1 H', J')$. A similar family of pairs induces a continuation map ϕ' from $SH_*^\#(\lambda H', J')$ to $SH_*^\#(\lambda Q_2 H, J)$. We also have

that $\phi' \circ \phi$ and $\phi \circ \phi'$ are directed system maps for $(SH_*^\#(\lambda H', J'))$ and $(SH_*^\#(\lambda H, J))$ respectively. This is because the directed system maps are also continuation maps induced by increasing families of Hamiltonians. Hence $(SH_*^\#(\lambda H, J))$ and $(SH_*^\#(\lambda H', J'))$ are isomorphic as directed systems. Also ϕ and its inverse ϕ' are restricted continuation morphisms. $\square$

Lemma 4.4. *Let $(H, J), (H', J')$ be growth rate admissible pairs such that $H' = H$ in a neighbourhood of $H^{-1}(0)$ and $H^{-1}(0) = H'^{-1}(0)$. Then $(SH_*^\#(\lambda H, J))$ is isomorphic to $(SH_*^\#(\lambda H', J'))$ as filtered directed systems. The isomorphism between these filtered directed systems is a composition of restricted continuation morphisms and switch morphisms and inverses of these morphisms.*

This Lemma is similar to Lemma 4.3 with one constraint removed.

Proof. of Lemma 4.4. From Lemma 4.2 there is a pair (H^p, J^p) constructed from H . This only depended on what H was on a small neighbourhood of $H^{-1}(0)$, so the equivalent construction (H'^p, J'^p) is equal to (H^p, J^p) . Hence all we need to do in this section is to prove that the filtered directed system $(SH_*^\#(\lambda H, J))$ is isomorphic to $(SH_*^\#(\lambda H^p, J^p))$.

There is a compact co-dimension 0 submanifold K' with boundary containing $H^{-1}(0) = (H^p)^{-1}(0)$ and all the 1-periodic orbits of λH^p for every $\lambda \in (0, \infty) \setminus A$ (where A is discrete). We can assume that K' is a subset of $U_1^H = \cap_\lambda U_\lambda^H$. It also has the property that any solution of the perturbed Cauchy-Riemann equations with respect to (H^p, J^p) with boundary in K' is contained in K' . Let $L : M \rightarrow \mathbb{R}$ be a Hamiltonian such that: $L = H^p$ on a small neighbourhood of K' and such that $L > 0$ outside K' . We also require that $L = H$ outside a closed subset of U_1^H . We let J_L be an almost complex structure compatible with the symplectic form such that it is equal to J^p on a small neighbourhood of K' and equal to J outside a closed subset of U_1^H . Because $L = H$ on a small neighbourhood of $H^{-1}(0) = L^{-1}(0)$ and $(L, J_L) = (H, J)$ outside a closed subset of U_1^H , we have that (L, J_L) is growth rate admissible. Let θ be the 1-form such that $\theta - \theta_M$ is exact and $-\theta(X_H) - H$ is bounded. Let $c > 0$ be a constant greater than the function $-\theta(X_L) - L + 1$. Let $L' : M \rightarrow \mathbb{R}$ be a function which is equal to L on a small neighbourhood of $L^{-1}(0)$ and equal to $L + c$ on a small neighbourhood of $\overline{M \setminus K'}$ and is greater than 0 everywhere else. Because $-\theta(X_{L'}) - L' < 0$ outside the interior of K' and no λ -periodic orbits of L' intersect the boundary of K' for λ outside some discrete set, we have that all the 1-periodic orbits of $\lambda L'$ of non-negative action are contained in the interior of K' . Hence by part 6 from Lemma 4.2, we have that $(\lambda L', J_L)$ satisfies the maximum principle with respect to $(K')^\circ$ (the interior of K') for all λ outside a discrete subset. So we can define $U_\lambda^{L'} := (K')^\circ$.

Let $\tilde{L}'$ be equal to L' inside K' and equal to $H^p + c$ outside K' . Again for all λ outside a discrete subset, all the 1-periodic orbits of non-negative action are contained in the interior K' . Hence part (6) from Lemma 4.2, ensures

that any compact curve satisfying Floer's equations with respect to $(\tilde{L}', J_L)$ whose boundary is contained in the interior of K' must be contained in K' . Hence $(\tilde{L}', J_L)$ satisfies the maximum principle with respect to $U_{\lambda}^{\tilde{L}'} := (K')^o$. Because $U_{\lambda}^{L'} = U_{\lambda}^{\tilde{L}'} = (K')^o$ for all λ and $(L', J_L) = (\tilde{L}', J_L)$ on a small neighbourhood of K' , we have a switch isomorphism from $(SH_{*}^{\#}(\lambda L', J_L))$ to $(SH_{*}^{\#}(\lambda \tilde{L}', J^p))$.

Lemma 4.3 tells us that $(SH_{*}^{\#}(\lambda \tilde{L}', J^p))$ and $(SH_{*}^{\#}(\lambda H^p, J^p))$ are isomorphic by a restricted continuation morphism. Lemma 4.3 also tells us that $(SH_{*}^{\#}(\lambda L', J))$ and $(SH_{*}^{\#}(\lambda H, J))$ are isomorphic. Hence $(SH_{*}^{\#}(\lambda H, J))$ and $(SH_{*}^{\#}(\lambda H^p, J^p))$ are isomorphic. $\square$

If $(H_1, J_1), (H_2, J_2)$ are growth rate admissible such that $H_1^{-1}(0)$ contains $H_2^{-1}(0)$ then there is a natural morphism of filtered directed systems from $(SH_{*}^{\#}(\lambda H_1, J_1))$ to $(SH_{*}^{\#}(\lambda H_2, J_2))$. Here is how this morphism is constructed: we have a non-decreasing family of Hamiltonians H^s with $H^s = H_1$ for $s \ll 0$ and such that $H^s = K$ for $s \gg 0$ where K satisfies:

- (1) $K^{-1}(0) = H_2^{-1}(0)$ and $H_2 = K$ on a small neighbourhood of $K^{-1}(0)$.
- (2) $K = H_1$ outside a closed subset of $U_1^{H_1}$.
- (3) (K, J_1) is growth rate admissible.

We also require that $H^s = H_1$ outside a closed subset of $U_1^{H_1}$. The family of pairs (H^s, J_1) induces a continuation map from $SH_{*}^{\#}(\lambda H_1, J_1)$ to $SH_{*}^{\#}(\lambda K, J_1)$. This in turn induces a morphism of filtered directed systems from $(SH_{*}^{\#}(\lambda H_1, J_1))$ to $(SH_{*}^{\#}(\lambda K, J_1))$ as the continuation map commutes with the filtered directed system maps. Our morphism is constructed by composing the above morphism with the isomorphism from $(SH_{*}^{\#}(\lambda K, J_1))$ to $(SH_{*}^{\#}(\lambda H_2, J_2))$ from Lemma 4.4. We call such a morphism a *growth rate admissible morphism*.

Lemma 4.5. *The composition of two growth rate admissible morphisms is a growth rate admissible morphism.*

Proof. of Lemma 4.5. The point is that composing switch morphisms gives us switch morphisms, composing restricted continuation maps gives us restricted continuation maps and if we have restricted continuation maps of the right form (so that some maximum principle still applies) then restricted continuation maps can commute with switch morphisms. Let $(H_1, J_1), (H_2, J_2), (H_3, J_3)$ be growth rate admissible pairs with $H_{i+1}^{-1}(0)$ contained in $H_i^{-1}(0)$. We wish to show that the composition of the growth rate admissible morphisms

$$E_1 : (SH_{*}^{\#}(\lambda H_1, J_1)) \rightarrow (SH_{*}^{\#}(\lambda H_2, J_2))$$

$$E_2 : (SH_{*}^{\#}(\lambda H_2, J_2)) \rightarrow (SH_{*}^{\#}(\lambda H_3, J_3))$$

is a growth rate admissible morphism

$$E_3 : SH_{*}^{\#}(\lambda H_1, J_1) \rightarrow SH_{*}^{\#}(\lambda H_3, J_3).$$

Let (H, J) be any growth rate admissible pair. By using methods from the proof of Lemma 4.4, we can construct Hamiltonians L_H^i and almost complex structures J_H^i for $i = 1, 2, 3$ with the following properties:

- (1) $(L_H^i)^{-1}(0) = H_i^{-1}(0)$ and $H_i = L_H^i$ near $H_i^{-1}(0)$.
- (2) $-\theta_M(X_{L_H^i}) - L_H^i \ll 0$ outside a small neighbourhood K_i of $H_i^{-1}(0)$.
- (3) Any map $s : S \rightarrow M$ with boundary inside K_i satisfying Floer's equation with respect to (L_H^i, J_H^i) must be contained in K_i . This statement is true even if we change (L_H^i, J_H^i) so that it equal to something else outside K_i as long as the Hamiltonian stays positive.
- (4) $K_3 \subset K_2 \subset K_1$.
- (5) (L_H^2, J_H^2) is equal to (L_H^3, J_H^3) outside K_3 and near ∂K_3 and (L_H^1, J_H^1) is equal to (L_H^2, J_H^2) outside K_2 and near ∂K_2 .
- (6) (L_H^i, J_H^i) is equal to $(H + \kappa, J)$ for some large constant $\kappa \gg 0$ near infinity.

Note that the pair (L_H^i, J_H^i) really depends on both H and J but we suppressed J from the notation to make it less cluttered. We now construct a smooth non-decreasing 1-parameter family of pairs (L_H^t, J_H^t) for $t \in [1, 3]$ such that

- (1) (L_H^i, J_H^i) are equal to the pairs constructed above with the same name for $i = 1, 2, 3$.
- (2) $(L_H^t, J_H^t) = (L_H^3, J_H^3)$ outside K_3 and near ∂K_3 .
- (3) For $t \in [1, 2]$, $(L_H^t, J_H^t) = (L_H^2, J_H^2)$ outside K_2 and near ∂K_2 .

This family gives us continuation maps:

$$\Psi_{i,j}^H : SH_*^\#(\lambda L_H^i, J_H^i) \rightarrow SH^+(\lambda L_H^j, J_H^j)$$

that do not depend on the choice of (H, J) for any $i \leq j, i, j = 1$ or 2 or 3 because no Floer trajectories connecting non-negative action orbits escapes the region K_3 . The point is that if we change (H, J) , we only do this outside K_3 . The map $\Psi_{i,j}^H$ induces a morphism of filtered directed systems from $(SH_*^\#(\lambda L_H^i, J_H^i))$ to $(SH^+(\lambda L_H^j, J_H^j))$. The filtered directed systems are canonically isomorphic for any choice of (H, J) via a switch morphism and the maps $\Psi_{i,j}^H$ are exactly the same for any choice of (H, J) . Hence we have filtered directed system maps $A_{i,j}^k$ that are isomorphisms from $(SH_*^\#(\lambda L_{H_i}^k, J_{H_i}^k))$ to $(SH^+(\lambda L_{H_j}^k, J_{H_j}^k))$. And we also have the relations

$$(2) \quad A_{k,k'}^j \circ \Psi_{i,j}^{H_k} \circ (A_{k,k'}^i)^{-1} = \Psi_{i,j}^{H_{k'}}.$$

We also have a restricted continuation morphism Φ_i of from the filtered directed system $SH_*^\#(\lambda L_{H_i}^i, J_{H_i}^i)$ to $SH_*^\#(\lambda H_i, J_i)$ (see Lemma 4.3). Its inverse Φ_i^{-1} is also a restricted continuation morphism. We have that $\Psi_{2,3}^{H_2} \circ \Psi_{1,2}^{H_2}$ is induced entirely from continuation maps and so it is exactly the same as the map $\Psi_{1,3}^{H_2}$. Also composing Φ_i or Φ_i^{-1} with any of these maps (when

possible) also induces restricted continuation morphisms. We get that the growth rate admissible morphism E_1 is equal to

$$\Phi_2 \circ A_{1,2}^2 \circ \Psi_{1,2}^{H_1} \circ \Phi_1^{-1}$$

and E_2 is equal to

$$\Phi_3 \circ A_{2,3}^3 \circ \Psi_{2,3}^{H_2} \circ \Phi_2^{-1}$$

by looking at the proof of Lemma 4.4 and the definition of growth rate admissible morphism. Hence their composition $E_2 \circ E_1$ is equal to:

$$\Phi_3 \circ A_{2,3}^3 \circ \Psi_{2,3}^{H_2} \circ A_{1,2}^2 \circ \Psi_{1,2}^{H_1} \circ \Phi_1^{-1}.$$

By equation 2 we have that this composition is equal to

$$\Phi_3 \circ A_{2,3}^3 \circ A_{1,2}^3 \circ \Psi_{2,3}^{H_1} \circ \Psi_{1,2}^{H_1} \circ \Phi_1^{-1} =$$

$$\Phi_3 \circ A_{2,3}^3 \circ A_{1,2}^3 \circ \Psi_{1,3}^{H_1} \circ \Phi_1^{-1}.$$

The composition of the switch morphisms $A_{2,3}^3 \circ A_{1,2}^3$ is the switch morphism $A_{1,3}^3$ hence we have that $E_2 \circ E_1$ is equal to:

$$\Phi_3 \circ A_{1,3}^3 \circ \Psi_{1,3}^{H_1} \circ \Phi_1^{-1}$$

which is equal to E_3 . Hence $E_2 \circ E_1 = E_3$ which gives us functoriality. $\square$

Lemma 4.6. *Let (H_s, J_s) be a smooth family of growth rate admissible pairs parameterized by $s \in \mathbb{R}$ such that for $s_1 < s_2$, $H_{s_1}^{-1}(0)$ contains $H_{s_2}^{-1}(0)$. For s_2 greater than s_1 , the growth rate admissible morphism from $(SH_*^\#(\lambda H_{s_1}, J_{s_1}))$ to $(SH_*^\#(\lambda H_{s_2}, J_{s_2}))$ is an isomorphism.*

Proof. of Lemma 4.6. By a compactness argument and by functoriality (Lemma 4.5) we only need to show that for each $s \in \mathbb{R}$, there is a $\delta_s > 0$ such that the result is true for $s_1 > s - \delta_s$ and $s_2 < s + \delta_s$. We will prove this by changing our pairs (H_s, J_s) to ones similar to the one described in Lemma 4.2. Then we note that if all the orbits have non-negative action then SH_* is equal to $SH_*^\#$ hence we can construct an inverse to our growth rate admissible morphism by using a decreasing family of Hamiltonians.

Fix the Hamiltonian H_s . We can ensure that there is a continuous family of small neighbourhoods $U_{s'}$ of $H_{s'}^{-1}(0)$ in which $-\theta(X_{H_{s'}}) - H_{s'} > 0$ on $U_{s'} \setminus H_{s'}^{-1}(0)$ with respect to s' by a 1-parameter version of the note after Lemma 4.2. This means that there is a $\delta_s > 0$ and a neighbourhood U of $H_s^{-1}(0)$ with $-\theta(X_{H_{s'}}) - H_{s'} > 0$ on $U \setminus H_{s'}^{-1}(0)$ for all $|s - s'| < \delta_s$.

We first perturb $H_{s-\delta_s}$ by a C^∞ small amount near its zero set to K so that $(K, J_{s-\delta_s})$ is still growth rate admissible and so that the interior of $K^{-1}(0)$ contains $H_{s-\delta_s}^{-1}(0)$ and $K^{-1}(0)$ is contained in U . By Lemma 4.2 let $(K^p, J_{s-\delta_s}^p)$ be a pair such that

- (1) $(K^p, J_{s-\delta_s}^p)$ is growth rate admissible.
- (2) $K^p = K$ on a small neighbourhood of $K^{-1}(0)$ and $(K^p)^{-1}(0) = K^{-1}(0)$.
- (3) $-\theta(K^p) - K^p \geq 0$ everywhere and $-\theta(K^p) - K^p > 0$ when $K^p > 0$.

We shrink U so that it is also contained in U_1^K . Choose an open set U' whose closure is contained in U and which still contains $K^{-1}(0)$. Let $\rho : M \rightarrow \mathbb{R}$ be a bump function which is equal to 1 inside U' and equal to 0 outside U . For $\kappa > 0$ large enough and for $|s - s'| \leq \delta_s$ we have that:

$$H'_{s'} := \rho H_{s'} + \kappa K^p$$

satisfies $-\theta(H'_{s'}) - H'_{s'} \geq 0$ because $-\theta(K^p) - K^p > 0$ in the relatively compact region $U \setminus U'$ and so is bounded below by a positive constant. We define a new family of almost complex structures $J'_{s'}$ to be equal to $J_{s-\delta_s}^p$ outside U' and equal to $J_{s'}$ on a small neighbourhood of $H_{s'}^{-1}(0)$. We have that $(H'_{s'}, J'_{s'})$ is growth rate admissible.

We have that $SH_*(\lambda H'_{s'}, J'_{s'})$ is a filtered directed system isomorphic to $SH_*^\#(\lambda H'_{s'}, J'_{s'})$ because $-\theta(H'_{s'}) - H'_{s'} \geq 0$. Also we have that if (K', Y') is any pair equal to $(H'_{s'}, J'_{s'})$ outside U' then $SH_*(\lambda K', Y')$ is equal to $SH_*(\lambda H'_{s'}, J'_{s'})$ as the maximum principle ensures that continuation maps between these Hamiltonians are well defined (and that these continuation maps do not have to be from non-decreasing families of Hamiltonians as we are not considering action). Hence for all $|s' - s| \leq \delta_s$, the filtered directed systems $(SH_*(\lambda H'_{s'}, J'_{s'}))$ are all isomorphic.

Let s_1, s_2 be such that $s - \delta_s < s_1 < s_2 < s + \delta_s$. We have a morphism ϕ of filtered directed systems from $(SH_*^\#(\lambda H'_{s_1}, J'_{s_1}))$ to $(SH_*^\#(\lambda H'_{s_2}, J'_{s_2}))$. By the previous discussion we also have a morphism ϕ' from

$$(SH_*^\#(\lambda H'_{s_2}, J'_{s_2})) \cong (SH_*(\lambda H'_{s_2}, J'_{s_2}))$$

to

$$(SH_*^\#(\lambda H'_{s_1}, J'_{s_1})) \cong (SH_*(\lambda H'_{s_1}, J'_{s_1})).$$

Because these morphisms are induced by continuation maps, we have that $\phi \circ \phi'$ and $\phi' \circ \phi$ are filtered directed system maps and hence by definition ϕ is an isomorphism of filtered directed systems.

By Lemma 4.4, we have an isomorphism Φ_i of filtered directed systems from $(SH_*^\#(\lambda H_{s_i}, J_{s_i}))$ to $(SH_*^\#(\lambda H'_{s_i}, J'_{s_i}))$ for $i = 1, 2$. This isomorphism and its inverse are growth rate admissible morphisms. We have an isomorphism $\Phi_2^{-1} \circ \phi \circ \Phi_1$ from $(SH_*^\#(\lambda H_{s_1}, J_{s_1}))$ to $(SH_*^\#(\lambda H_{s_2}, J_{s_2}))$. Because $\phi, \Phi_1, \Phi_2^{-1}$ are growth rate admissible morphisms, we have by functoriality (Lemma 4.5) that $\Phi_2^{-1} \circ \phi \circ \Phi_1$ is also a growth rate admissible morphism. Hence the natural morphism from $(SH_*^\#(\lambda H_{s_1}, J_{s_1}))$ to $(SH_*^\#(\lambda H_{s_2}, J_{s_2}))$ is an isomorphism. $\square$

Lemma 4.7. *Let $(H_0, J_0), (H_1, J_1)$ be two growth rate admissible Hamiltonians such that $H_1^{-1}(0) \subset H_0^{-1}(0)$. Suppose that there is a smooth family of Hamiltonians G_t satisfying the Liouville vector field property such that*

- (1) $G_{s_2}^{-1}(0) \subset G_{s_1}^{-1}(0)$ if $s_1 \leq s_2$.
- (2) $G_0 = H_0$ on a neighbourhood of $H_0^{-1}(0)$ and $G_0^{-1}(0) = H_0^{-1}(0)$.
- (3) $G_1 = H_1$ on a neighbourhood of $H_1^{-1}(0)$ and $G_1^{-1}(0) = H_1^{-1}(0)$.

Then the growth rate admissible morphism

$$(SH_*^\#(\lambda H_0, J_0)) \rightarrow (SH_*^\#(\lambda H_1, J_1))$$

is an isomorphism.

Proof. of Lemma 4.7. By using construction similar to the one in the proof of Lemma 4.6 combined with Lemma 4.2, we have for each $s \in [0, 1]$ there is a $\delta_s > 0$ and a smooth family of growth rate admissible pairs (G_s^t, J_s^t) ($t \in (s - \delta_s, s + \delta_s)$) satisfying:

- (1) $G_s^t = G_t$ on a neighbourhood of $G_t^{-1}(0)$ and $G_t^{-1}(0) = (G_s^t)^{-1}(0)$.
- (2) $(G_0^0, J_0^0) = (H_0, J_0)$ on a neighbourhood of $H_0^{-1}(0)$ and $(G_0^0)^{-1}(0) = H_0^{-1}(0)$.
- (3) $(G_1^1, J_1^1) = (H_1, J_1)$ on a neighbourhood of $H_1^{-1}(0)$ and $(G_1^1)^{-1}(0) = H_1^{-1}(0)$.

Hence by Lemma 4.6, we have for $s - \delta_s < t_1 \leq t_2 < t + \delta_s$ the natural filtered directed system map from $(SH_*^\#(G_s^{t_1}, J_s^{t_1}))$ to $(SH_*^\#(G_s^{t_2}, J_s^{t_2}))$ is an isomorphism.

Hence by a compactness argument there is a sequence of growth rate admissible pairs $(A_i, Y_i) := (G_{s_i}^{t_i}, J_{s_i}^{t_i})$ for $i = 1, \dots, k$ such that

- (1) $A_i^{-1}(0)$ contains $A_{i+1}^{-1}(0)$.
- (2) $(A_k, Y_k) = (G_1^1, J_1^1)$ and $(A_0, Y_0) = (G_0^0, J_0^0)$.
- (3) The morphism from $(SH_*^\#(A_i, Y_i))$ to $(SH_*^\#(A_{i+1}, Y_{i+1}))$ is an isomorphism.

We also have by Lemma 4.4 that the growth rate admissible morphisms

$$(SH_*^\#(H_0, J_0)) \rightarrow (SH_*^\#(A_0, Y_0))$$

and

$$(SH_*^\#(A_1, Y_1)) \rightarrow (SH_*^\#(H_1, J_1))$$

are isomorphisms. Hence by functoriality of these morphisms we get that the growth rate admissible morphism from $(SH_*^\#(\lambda H_0, J_0))$ to $(SH_*^\#(\lambda H_1, J_1))$ is an isomorphism. $\square$

Lemma 4.8. *Let (H, J) be a growth rate admissible pair. For any compact set $K \subset M$, there exists a growth rate admissible pair (H_K, J_K) such that:*

- (1) $H_K^{-1}(0)$ contains both $H^{-1}(0)$ and K .
- (2) The morphism from $(SH_*^\#(\lambda H_K, J_K))$ to $(SH_*^\#(\lambda H, J))$ is an isomorphism.

Proof. of Lemma 4.8. For the purposes of this proof we may as well enlarge K so that it contains $H^{-1}(0)$. By Lemma 4.7 all we need to do is create a family of Hamiltonians $H_s, (s \in [0, \infty])$ such that

- (1) H_s satisfies the Liouville vector field property.
- (2) $H_{s_2}^{-1}(0)$ contains $H_{s_1}^{-1}(0)$ if $s_1 \leq s_2$.
- (3) $H_0 = H$ and $H_s^{-1}(0)$ contains K for $s \gg 1$.

Let f_H, X_{θ_H} be the function and Liouville vector field which enables (H, J) to satisfy the Liouville vector field property. Let ρ be a bump function such that it is equal to 1 on a neighbourhood of K and 0 outside some larger compact set. Because the vector field $V'_H := \rho X_{\theta_H}$ has compact support, its flow $\phi_t^{V'_H}$ is well defined everywhere. Let $H_s := ((\phi_s^{V'_H})_* H)$. For $\epsilon_H > 0$ small enough we have that $dH(X_{\theta_H}) > 0$ inside $H^{-1}(0, \epsilon_H)$. Because $(\phi_t^{V'_H})^* X_{\theta_H}$ is proportional to X_{θ_H} we have that $dH_s(X_{\theta_H}) = (\phi_t^{V'_H})_* \left(dH((\phi_t^{V'_H})^* X_{\theta_H}) \right) > 0$ in the region $H_s^{-1}(0, \epsilon_H)$. Also because H satisfies the Liouville vector field property, there exists a C_s such that $f_H^{-1}(-\infty, C_s]$ is contained in $H_s^{-1}(0)$ and $df_H(X_{\theta_H}) > 0$ in $f_H^{-1}(C_s, \infty)$. Hence H_s satisfies the Liouville vector field property for all $s \geq 0$. There is a constant C so that $f_H^{-1}(-\infty, C] \subset H^{-1}(0)$ and $df_H(V'_H) > 0$ in the region $K \setminus f_H^{-1}(-\infty, C)$. Hence the time s flow of $f_H^{-1}(-\infty, C]$ contains K for $s \gg 1$. Because $f_H^{-1}(-\infty, C]$ is contained inside $H^{-1}(0)$ we have for $s \gg 1$ that $H_s^{-1}(0)$ contains K .

By Lemma 4.2 there is a growth rate admissible pair (H_K, J_K) such that $H_K^{-1}(0) = G_s^{-1}(0)$ for some $s \gg 0$ and such that $H_K = G_s$ on a neighbourhood of $H_K^{-1}(0)$. We assume that s is large enough so that $H_K^{-1}(0)$ contains K . Hence by Lemma 4.7 we have that the growth rate admissible morphism

$$(SH_*^\#(\lambda H_K, J_K)) \rightarrow (SH_*^\#(\lambda H, J))$$

is an isomorphism. $\square$

Proof. of Theorem 4.1. Let K_1 be a compact set whose interior contains both $H^{-1}(0)$ and $H'^{-1}(0)$. By Lemma 4.8, there is a pair (H_1, J_1) with $H_1^{-1}(0)$ containing K_1 such that the growth rate admissible morphism from $(SH_*^\#(\lambda H_1, J_1))$ to $(SH_*^\#(\lambda H, J))$ is an isomorphism. Choose a compact set K_2 whose interior contains $H_1^{-1}(0)$. Let (H'_1, J'_1) be a pair such that $(H'_1)^{-1}(0)$ contains K_2 and such that the morphism from $(SH_*^\#(\lambda H'_1, J'_1))$ to $(SH_*^\#(\lambda H', J'))$ is an isomorphism. By repeating this process, we can find two more growth rate admissible pairs $(H_2, J_2), (H'_2, J'_2)$ such that:

- (1) $(H'_1)^{-1}(0)$ is contained in the interior of $H_2^{-1}(0)$.
- (2) The morphism from $(SH_*^\#(\lambda H_2, J_2))$ to $(SH_*^\#(\lambda H_1, J_1))$ is an isomorphism.
- (3) $H_2^{-1}(0)$ is contained in the interior of $(H'_2)^{-1}(0)$.
- (4) The morphism from $(SH_*^\#(\lambda H'_2, J'_2))$ to $(SH_*^\#(\lambda H'_1, J'_1))$ is an isomorphism.

Hence we have the following sequence of morphisms:

$$\begin{aligned} (SH_*^\#(\lambda H'_2, J'_2)) &\rightarrow (SH_*^\#(\lambda H_2, J_2)) \rightarrow \\ &(SH_*^\#(\lambda H'_1, J'_1)) \rightarrow (SH_*^\#(\lambda H_1, J_1)) \end{aligned}$$

where composing any two of these morphisms gives an isomorphism. By Theorem 3.2, the middle morphism from the group $(SH_*^\#(\lambda H_2, J_2))$ to

$(SH_*^\#(\lambda H'_1, J'_1))$ is an isomorphism. Hence $(SH_*^\#(\lambda H, J))$ is isomorphic to $(SH_*^\#(\lambda H', J'))$. This proves the theorem. $\square$

Corollary 4.9. *The filtered directed system $(SH_*^\#(\lambda H, J))$ is an invariant of M up to exact symplectomorphism as long as this symplectomorphism preserves our choice of trivialization τ of the canonical bundle and our class $b \in H^2(M, \mathbb{Z}/2\mathbb{Z})$. If M is complete (i.e. the ω_M -dual of θ_M is an integrable vector field) then it is an invariant up to general symplectomorphism (again preserving (τ, b)).*

Proof. All the properties defining growth rate admissibility, are invariants of M up to exact symplectomorphism, hence by Theorem 4.1 we have that $(SH_*^\#(\lambda M, J))$ is an invariant up to exact symplectomorphism preserving (τ, b) . Suppose that M is complete. Then it is the completion of some Liouville domain $\overline{M}$ hence by [BTE09, Lemma 1], if M, M' are symplectomorphic then they are exact symplectomorphic. Hence $(SH_*^\#(\lambda M, J))$ is an invariant up to general symplectomorphism preserving (τ, b) in this case. $\square$

Because its an invariant up to exact symplectomorphism preserving (τ, b) , we will write:

$$(SH_*^\#(M, \theta, \lambda))$$

for any filtered directed system $(SH_*^\#(\lambda H, J))$ where we have chosen some growth rate admissible pair (H, J) and a pair (τ, b) .

If (N, θ_N) is a Liouville domain, then the interior of N , N^0 is a finite type convex symplectic manifold for the following reason: Let X_{θ_N} be the $d\theta_N$ -dual of θ_N . By flowing back ∂N backwards along X_{θ_N} , we get that a collar neighbourhood of ∂N is equal to $(1 - \epsilon, 1] \times \partial N$ with $\theta_N = r_N \alpha_N$. Here r_N parameterizes the interval and $\alpha_N = \theta_N|_{\partial N}$. Let $g : (1 - \epsilon, 1) \rightarrow \mathbb{R}$ be a function which is equal to 0 near $1 - \epsilon$ and tends to $+\infty$ near 1 and also that its derivative is positive near 1. We let $f_N : N^0 \rightarrow \mathbb{R}$ be a function which is 0 away from this collar neighbourhood and equal to $g(r_N)$ inside this collar neighbourhood. This shows that (N^0, θ_N) has the structure of a finite type convex symplectic manifold.

Lemma 4.10. *Let (M, θ) be a finite type convex symplectic manifold and let X_θ be the $d\theta$ -dual of θ . Let $f_M : M \rightarrow \mathbb{R}$ be the exhausting function such that $df_M(X_\theta) > 0$ in $f_M^{-1}[C, \infty)$ for some $C \gg 0$. We define N to be the Liouville domain $f_M^{-1}(-\infty, C]$. We also assume that the period spectrum of the contact manifold ∂N is discrete. Then the filtered directed systems $(SH_*(M, \theta, \lambda))$ and $(SH_*(N^0, \theta|_{N^0}, \lambda))$ are isomorphic as long as the choice of trivialization τ and homology class b for N^0 is equal to such a choice for M restricted to N^0 .*

Proof. of Lemma 4.10. Let $U \subset [1, \infty) \times \partial N$ be the partial cylindrical end of M obtained by flowing ∂N along X_θ . Let r be the coordinate parameterizing $[1, \infty)$. We have that $\theta = r\alpha$ inside U where $\alpha = \theta|_{\partial N}$. By flowing ∂N backwards along X_θ we can extend U to U' (containing $\overline{M \setminus N}$) so that it

is now a subset of $(0, \infty) \times \partial N$. We also extend r so it now parameterizes the larger interval $(0, \infty)$. Let $h : (0, \infty) \rightarrow \mathbb{R}$ be a function such that $h(r) = 0$ near $r = 0$ and $h(r) = r$ for $r \geq 1 - \delta$ where $\delta > 0$ is small. We also assume that $h \geq 0$ and $h' > 0$ for $h > 0$. We set H to be equal to $h(r)$ where r is well defined and 0 otherwise. We let J be an almost complex structure on M such that it is cylindrical on the region $\{r \geq 1 - \delta\}$. The pair (H, J) is growth rate admissible (this is because it is basically the same as the pair from the second example mentioned earlier). The pair $(H|_{N^0}, J|_{N^0})$ is also growth rate admissible. Let $A \subset (0, \infty)$ be the period spectrum of ∂N . If $\lambda \in (0, \infty) \setminus A$ then all the 1-periodic orbits of λH are contained in N^0 . Also because J is cylindrical, by Corollary 9.3 all the Floer trajectories connecting orbits of λH or continuation map Floer trajectories joining $\lambda_1 H$ and $\lambda_2 H$ are contained in N^0 . This ensures that the filtered directed systems $(SH_*^\#(\lambda H, J))$ and $(SH_*^\#(\lambda H|_{N^0}, J|_{N^0}))$ are isomorphic. This completes the Lemma. $\square$

Lemma 4.11. *Suppose that (M, θ) and (M', θ') are convex deformation equivalent, then $(SH_*^\#(M, \theta, \lambda))$ and $(SH_*^\#(M', \theta', \lambda))$ are isomorphic as filtered directed systems (again the choice of trivialization τ and homology class b for M' must be the same as that of M).*

Proof. of Lemma 4.11. Let f_M, X_θ, N be as in the previous Lemma. By Corollary 8.3, we have that (M, θ) is convex deformation equivalent to the completion $(\widehat{N}, \theta_N)$. Let r_N be the cylindrical coordinate of $\widehat{N}$. We can extend the cylindrical end $[1, \infty) \times \partial N$ to $(0, \infty) \times \partial N$ inside $\widehat{N}$ by flowing ∂N backwards along the Liouville vector field X_{θ_N} . By applying Lemma 4.10 twice (once to (M, θ) and once to $(\widehat{N}, \theta_N)$) we get that $(SH_*^\#(M, \theta, \lambda))$ is isomorphic to $(SH_*^\#(N^0, \theta|_N = \theta_N|_N, \lambda))$ which is isomorphic to $(SH_*^\#(\widehat{N}, \theta_N, \lambda))$.

Similarly we have that there is a Liouville domain N' such that the filtered directed system $(SH_*^\#(M', \theta', \lambda))$ is isomorphic to $(SH_*^\#(\widehat{N}', \theta_{N'}, \lambda))$ and such that (M', θ') is convex deformation equivalent to $(\widehat{N}', \theta_{N'})$. Because convex deformation equivalence is an equivalence relation we get that $(\widehat{N}, \theta_N)$ is convex deformation equivalent to $(\widehat{N}', \theta_{N'})$. Also both $(\widehat{N}, \theta_N)$ and $(\widehat{N}', \theta_{N'})$ are complete so by Corollary 8.6 they are exact symplectomorphic. Hence by Lemma 4.9 we have that the groups $(SH_*^\#(\widehat{N}, \theta_N, \lambda))$ and $(SH_*^\#(\widehat{N}', \theta_{N'}, \lambda))$ are isomorphic. This implies that $(SH_*(M, \theta, \lambda))$ and $(SH_*(M', \theta', \lambda))$ are isomorphic. $\square$

Motivated by Lemma 4.9 we have the following definition:

Definition 4.12. *For any finite type convex symplectic manifold M , we define*

$$\Gamma(M, \theta) := \Gamma((SH_*^\#(M, \theta, \lambda))).$$

Sometimes we write $\Gamma(M)$ if it is clear what the Liouville form θ is.

We will show later in Corollary 4.15 that this is the same as growth rate as in Definition 2.3.

In some cases, we wish to consider orbits of all actions and not just ones of non-negative action. Let (H, J) be growth rate admissible such that $-\theta_H(X_H) - H \geq 0$ for some θ_H where $\theta_H - \theta_M$ is exact. Let (H_λ, J_λ) be a smooth family of Hamiltonians parameterized by $\lambda \geq 1$ such that $(H_\lambda, J_\lambda) = (\lambda H + c_\lambda, J)$ outside a closed subset of U_λ^H . Here c_λ is a smooth family of constants. Basically by the maximum principle we have that $SH_*(H_\lambda, J_\lambda)$ is well defined and for $\lambda_1 < \lambda_2$ there is a morphism from $SH_*(H_{\lambda_1}, J_{\lambda_1})$ to $SH_*(H_{\lambda_2}, J_{\lambda_2})$. Note that outside a closed subset of U_λ^H , λH has no 1-periodic orbits of negative action and hence no 1-periodic orbits outside this closed subset. This morphism is induced by the smooth family of pairs (H_λ, J_λ) from λ_1 to λ_2 . Hence $(SH_*(H_\lambda, J_\lambda))$ forms a filtered directed system.

Lemma 4.13. *Suppose that $-\theta(X_H) - H \geq 0$ then $(SH_*(H_\lambda, J_\lambda))$ is isomorphic to $(SH_*^\#(M, \theta, \lambda))$.*

Proof. of Lemma 4.13. Because the action of all the 1-periodic orbits of λH are non-negative, we have that $SH_*(\lambda H, J) \cong SH_*^\#(\lambda H, J)$. This isomorphism commutes with the filtered directed system maps because they are continuation maps induced from an increasing family of Hamiltonians. Hence the filtered directed system $(SH_*(\lambda H, J))$ is isomorphic to $(SH_*^\#(\lambda H, J))$.

Let $q : \mathbb{R} \rightarrow [0, 1]$ be a smooth function such that $q(x) = 0$ for $x \leq 0$ and $q(x) = 1$ for $x \geq 1$. By joining $(\lambda H, J)$ with (H_λ, J_λ) via a smooth family of pairs $(H_\lambda^s, J_\lambda^s)$ such that $(H_\lambda^s, J_\lambda^s) = (\lambda H + q(s)c_\lambda, J)$ outside a closed subset of U_λ^H , we have by the maximum principle a well defined continuation isomorphism from $SH_*(\lambda H, J)$ to $SH_*(H_\lambda, J_\lambda)$. This monomorphism commutes with the continuation maps. Also the continuation map induced by the family $(H_\lambda^{-s}, J_\lambda^{-s})$ gives us an inverse to the above morphism. This is because the composition of these two continuation maps is a continuation map induced by some family of pairs equal to $(\lambda H + q(s)c'_\lambda, J)$ near infinity and these are homotopic through such families of pairs to the constant pair. The constant pair gives us the identity map.

Hence we have that the filtered directed system $(SH_*(\lambda H, J))$ is isomorphic to $(SH_*(H_\lambda, J_\lambda))$. Hence we have that $(SH_*^\#(\lambda H, J))$ is isomorphic to $(SH_*(H_\lambda, J_\lambda))$. This proves the Lemma. $\square$

Corollary 4.14. *Let N be a Liouville domain and let r_N be the cylindrical coordinate of $\hat{N}$. Suppose also that the period spectrum of ∂N is discrete. Let J be an almost complex structure that is cylindrical near infinity. Let (H_λ, J_λ) be a family of Hamiltonians such that $H_\lambda = \lambda r_N + c_\lambda$ outside a large compact set K for some family of constants c_λ . Suppose also that J_λ is equal to J outside K . Then $(SH_*(H_\lambda, J_\lambda))$ is isomorphic as a filtered directed system to $(SH_*^\#(\hat{N}, \theta_N, \lambda))$.*

Proof. of Corollary 4.14. Let $h : [1, \infty) \rightarrow \mathbb{R}$ be a function such that $h(x) = 0$ for x near 1 and $h(x) = x - 2$ for $x \geq 3$. Suppose also that $h'(x), h''(x) \geq 0$. Let H be a Hamiltonian such that $H = 0$ inside $N \subset \hat{N}$ and $H = h(r_N)$ outside N . Inside N , $-\theta(X_H) - H = 0$. Outside N ,

$$-\theta(X_H) - H = r_N h'(r_N) - h(r_N) = \int_1^{r_N} x h''(x) dx \geq 0.$$

Hence $-\theta(X_H) - H \geq 0$. This pair is growth rate admissible for reasons similar to the reason why the second example mentioned earlier is growth rate admissible. Hence by Lemma 4.13 we get our result. $\square$

The problem is that the growth rate $\Gamma(M, b)$ has a different definition to the one given in definition 2.3. We recall the definition here: In section 2.2, we defined $SH_*^{\leq \lambda}(N)$ for a Liouville domain N which is the direct limit of $SH_*^{\leq \lambda}(H, J)$ where (H, J) is a pair defined on $\hat{N}$ and is cylindrical at infinity and less than 0 on N . For $\lambda_1 \leq \lambda_2$, there is a natural map $SH_*^{\leq \lambda_1}(N) \rightarrow SH_*^{\leq \lambda_2}(N)$. This is a filtered directed system $(SH_*^{\leq \lambda}(N))$ whose direct limit is $SH_*(N)$.


Lemma 4.15. *The filtered directed system $(SH_*^{\leq \lambda}(N))$ is isomorphic to $(SH_*(\hat{N}, \theta_N, \lambda))$. Hence definitions 2.3 and 4.12 are equivalent. We assume that the period spectrum on N is discrete.*

Proof. of Lemma 4.15. Let (H, J) be the pair defined in the proof of Corollary 4.14. We will first construct a family of pairs (H_λ, J_λ) . Let r be the cylindrical coordinate of N . We will construct an isomorphism from $SH_*(\lambda H, J)$ to $SH_*^{\leq \lambda}(N)$ that commutes with the filtered directed system maps.

Fix $\lambda \geq 1$. We construct the Hamiltonian H_a as follows: H_a is constant and equal to $-\frac{1}{a}$ inside N . This construction only works if a is sufficiently large. In the region $\hat{N} \setminus N$ we let $H_a = g_a(r)$ where $g_a(r) = -\frac{1}{a}$ near $r = 1$, $g'_a(r), g''_a(r) \geq 0$ and $g_a(r) = a(r - 1 - \frac{1}{a})$ for $r \geq 1 + \frac{1}{a}$. We also require that in a neighbourhood of the region $1 + \frac{2}{3a\lambda} \leq r \leq 1 + \frac{5}{6a\lambda}$ that $g(a)$ is equal to

$$-\frac{1}{a} + (\lambda - \frac{3}{2a})(r - 1 - \frac{1}{2a\lambda - 3}).$$

Here is a picture:



We let J_a be anything we like inside N and J_a is cylindrical outside N . If we have a periodic orbit inside the level set $r = c$, then its action is $cg'(c) - g(c)$. The derivative of this with respect to c is $g'(c) + cg''(c) - g'(c) = cg''(c)$. Because $g'' \geq 0$ we have that orbits in the level set $r = c_1$ have action less than or equal to the ones in the level set $r = c_2$ for $c_1 \leq c_2$. Let $e = 1 + \frac{2}{3a\lambda}$. All the 1-periodic orbits in the region $r \leq e$ have action less than or equal to $eg'(e) - g(e)$ and all the orbits in the region $r \geq e$ have action greater than or equal to this same quantity. Also because $g(r) = -\frac{1}{a} + (\lambda - \frac{3}{2a})(r - 1 - \frac{1}{2a\lambda - 3})$ in the region $(e - \epsilon, e]$ for $\epsilon > 0$ small,

$$\begin{aligned}
 eg'(e) - g(e) &= e(\lambda - \frac{3}{2a}) + \frac{1}{a} - (\lambda - \frac{3}{2a})(e - 1 - \frac{1}{2a\lambda - 3}) \\
 &= \frac{1}{a} + (\lambda - \frac{3}{2a})(1 + \frac{1}{2a\lambda - 3}) \\
 &= \frac{1}{a} + (\lambda - \frac{3}{2a})(1 + \frac{\frac{1}{2a}}{\lambda - \frac{3}{2a}}) \\
 &= \frac{1}{a} + \lambda - \frac{3}{2a} + \frac{1}{2a} = \lambda.
 \end{aligned}$$

Because J is cylindrical and H is linear in the region $(e - \epsilon, e]$, we can use the maximum principle Corollary 9.3 to ensure that any Floer trajectory connecting orbits of action $\leq \lambda$ must be contained in $\{r \leq e\}$.

We can construct another growth rate admissible pair (H'_a, J'_a) where $(H'_a, J'_a) = (H_a, J_a)$ in the region $\{r \leq e\}$ and $H'_a = -\frac{1}{a} + (\lambda - \frac{3}{2a})(r - 1 - \frac{1}{2a\lambda - 3})$ outside this region. Then $SH_*(H'_a, J'_a) = SH_*^{\leq \lambda}(H_a, J_a)$. Also for

$a_1 \leq a_2$, a similar maximum principle argument ensures that the transfer map

$$SH_*(H'_{a_1}, J'_{a_1}) \rightarrow SH_*(H'_{a_2}, J'_{a_2})$$

coming from the family $(H'_{a_1+t(a_2-a_1)}, J'_{a_1+t(a_2-a_1)})$ is identical to the transfer map

$$SH_*^{\leq \lambda}(H_{a_1}, J_{a_1}) \rightarrow SH_*^{\leq \lambda}(H_{a_2}, J_{a_2})$$

coming from the family $(H_{a_1+t(a_2-a_1)}, J_{a_1+t(a_2-a_1)})$.

Also because the slope of H'_a is equal to $\lambda - \frac{3}{2a}$ near infinity we have that $SH_*(\lambda H, J)$ is isomorphic to $\varinjlim_a SH_*(H'_a, J'_a)$. Hence $SH_*(\lambda H, J) \cong SH_*^{\leq \lambda}(N)$ because $\varinjlim_a SH_*(H'_a, J'_a)$ is equal to $\varinjlim_a SH_*^{\leq \lambda}(H_a, J_a)$. By ensuring that the pairs $(H_a, J_a), (H'_a, J'_a)$ smoothly increase with λ and looking at the resulting continuation maps (and by Corollary 9.3 we get that the filtered directed system maps for $(SH_*(\lambda H, J))$ are identical to the ones for $(SH_*^{\leq \lambda}(N))$). Hence these directed systems are isomorphic. Hence $(SH_*^\#(\hat{N}, \theta_N, \lambda))$ is isomorphic to $(SH_*^{\leq \lambda}(N))$ by Corollary 4.14. $\square$

We will now prove Theorem 2.4 which says that growth rate is an invariant of a finite type Liouville manifold up to symplectomorphism.

Proof. of Theorem 2.4. By Corollary 4.15 we have that growth rate is equal to $\Gamma(SH_*^\#(M, \theta_M, \lambda))$. By Lemmas 4.9 and 3.1, we have that the growth rate $\Gamma(SH_*^\#(M, \theta_M, \lambda))$ is an invariant of M up to symplectomorphism. Hence $\Gamma(M)$ is an invariant of M up to symplectomorphism preserving the class $b \in H^2(M, \mathbb{Z}/2\mathbb{Z})$ and the choice of trivialization τ . $\square$

4.2. Growth rate of cotangent bundles. We let $\mathbb{K}$ be a field. Let (Q, g) be a Riemannian manifold, and let $L^\lambda := \mathcal{L}^{\leq \lambda^2}(Q, g)$ be the space of free loops of length $\leq \lambda^2$. For $\lambda_1 \leq \lambda_2$ we have a natural inclusion $L^{\lambda_1} \hookrightarrow L^{\lambda_2}$. This gives us a filtered directed system $(H_*(L^\lambda, \mathbb{K}))$. We define $\Gamma(Q, \mathbb{K})$ to be equal to $\Gamma((H_*(L^\lambda, \mathbb{K})))$. Note that Q has exponential growth if and only if there is some field $\mathbb{K}$ such that $\Gamma(Q, \mathbb{K}) = \infty$. We define $\omega_2 \in H^2(T^*Q, \mathbb{Z}/2\mathbb{Z})$ to be the pullback of the second Stiefel-Whitney class of Q . We also have a canonical choice of trivialization τ_Q of the canonical bundle of T^*Q induced by the volume form on Q . The cotangent bundle T^*Q is the completion of the unit cotangent bundle D^*Q which is a Liouville domain. Let θ_Q be the Liouville form on T^*Q . This is locally equal to $\sum_i p_i dq_i$ where p_i are momentum coordinates and q_i are position coordinates in Q . In this section we will prove:

Theorem 4.16. *The filtered directed systems $(H_*(L^\lambda))$ and $(SH_*^\#(T^*Q, \theta_Q, \lambda, (\tau_Q, \omega_2)))$ are isomorphic.*

This means we get the following Corollary:

Corollary 4.17. $\Gamma(T^*Q, (\tau_Q, \omega_2)) = \Gamma(Q)$.

Before we prove this theorem we need a slightly different definition of growth rate. Let (N, θ_N) be a Liouville domain. Let r_N be the radial coordinate for the cylindrical end $\partial N \times [1, \infty)$ of $\widehat{N}$. We assume that the period spectrum of the contact boundary ∂N is a discrete subset $\mathcal{P}$ of $\mathbb{R}$. We say that a Hamiltonian $H : S^1 \times \widehat{N} \rightarrow \mathbb{R}$ is *quadratic admissible* if:

- (1) there exists constants $b, b' \in \mathbb{R}$ such that $H + b \leq \frac{1}{2}r_N^2 \leq H + b'$ in the region $r_N \geq 1$. We also require that: $r_N + b \leq dH(\frac{\partial}{\partial r_N}) \leq r_N + b'$. This ensures that if H has some 1-periodic orbit inside the region $\alpha \leq r_N \leq \beta$ then its action is greater than $\alpha(\alpha + b) + b' - \frac{1}{2}\beta^2$ and less than $\beta(\beta + b') + b - \frac{1}{2}\alpha^2$.
- (2) There is a sequence $l_1 < l_2 < \dots$ tending to infinity such that $l_i \notin \mathcal{P}$ and $H = \frac{1}{2}r_N^2 + d_i$ on a small neighbourhood of the hypersurface $\{r_N = l_i\}$ where d_i is a constant (in particular H has no 1-periodic orbits near these hypersurfaces). This also ensures that the 1-periodic orbits starting at say $r_N = \alpha$ do not stray too far away from this hypersurface.
- (3) the sequence l_i satisfies $l_i/l_{i-1} < \kappa$ where κ is a constant.
- (4) We need that $H \geq l_i r_N - \frac{1}{2}l_i^2 + d_i$ in the region $r_N \geq l_i$.

By Lemma 2.2 we can perturb any quadratic admissible Hamiltonian so it becomes non-degenerate and remains quadratic admissible. Let H be such a Hamiltonian. We also define an almost complex structure J such that J is equal to some cylindrical almost complex structure on a neighbourhood of $\{r_N = l_i\}$ for each i . We say that J is compatible with H if this is true. Such a pair (H, J) is said to be *quadratic admissible*. Lemma 9.1 tells us that we can define $SH_*^{\leq \lambda}(H, J)$ (see [Rit98, Section 21.3] for an alternative way of defining this). These groups form a filtered directed system where all the filtered directed system maps come from the natural inclusions. Hence we have a filtered directed system $(SH_*^{\leq \lambda^2}(H, J))$.

Lemma 4.18. *The filtered directed system $(SH_*^{\leq \lambda^2}(H, J))$ is isomorphic to $(SH_*^{\leq \lambda}(N))$.*

We need a preliminary algebraic lemma.

Lemma 4.19. *Let $(V_x), (V'_x)$ be two filtered directed systems and let $1 \leq l_1 < l_2 < \dots$ be a sequence tending to infinity such that $l_i/l_{i-1} < K$ for some constant K . Suppose for each $x \in [1, \infty)$ we have a map $p_{x,i}$ from V_x to V'_{l_i} whenever $l_i \geq cx$ where $c \geq 1$ is some constant. We assume that p_x commutes with the filtered directed system maps (i.e. if $a_{x,y} : V_x \rightarrow V_y$ and $a'_{i,j} : V'_{l_i} \rightarrow V'_{l_j}$ are filtered directed system maps then we require that $p_{y,j} \circ a_{x,y} = a'_{i,j} \circ a_{x,i}$). Suppose we also have maps $p'_{i,x} : V'_{l_i} \rightarrow V_x$ for $l_i \leq c'x$ which also commute with the filtered directed system maps (here $c' \geq 1$ is a constant). Then (V_x) is isomorphic to (V'_x) .*

Proof. of Lemma 4.19. Let $a_{x,y} : V_x \rightarrow V_y$ and $a'_{x,y} : V'_x \rightarrow V'_y$ be the directed system maps. We first construct a map $\phi : V_x \rightarrow V'_{Kx}$ as follows:

choose l_i so that $1 \leq l_i/x < K$. We define ϕ as $a'_{l_i, Kx} \circ p_{x,i}$. We can define $\phi' : V'_x \rightarrow V'_{Kx}$ in a similar way as $p'_{i, Kx} \circ a'_{x, l_i}$. Because $p_{x,i}, p'_{i,x}$ commutes with the directed system maps, we have that $\phi \circ \phi'$ and $\phi' \circ \phi$ are directed system maps and hence we have an isomorphism. $\square$

Proof. of Lemma 4.18. We have a sequence l_i satisfying $l_i/l_{i-1} < \kappa$ where $H = \frac{1}{2}r_N^2 + d_i$ in a neighbourhood of $r_N = l_i$. Here d_i is a constant. We define H_i as follows:

- (1) $H_i = H$ in the region $r_N \leq l_i - \delta_i$ where δ_i is a very small constant.
- (2) $H_i = f_i(r_N)$ on a small neighbourhood of $l_i - \delta_i \leq r_N \leq l_i$ where $f'_i, f''_i \geq 0$. We can then ensure that H_i has no orbits in this neighbourhood.
- (3) $H_i = l_i r_N - \frac{1}{2}l_i^2 + d_i$ in the region $r_N \geq l_i$.
- (4) $H \geq H_i$ for all i (this can be done by the last condition that a quadratic admissible Hamiltonian satisfies). We also want that $H_{i-1} \leq H_i$ for all i .

Let K_s , $s \geq 1$ be a smooth family of Hamiltonians such that:

- (1) $\frac{\partial K_s}{\partial s} \geq 0$.
- (2) K_s is linear at infinity of slope s .
- (3) $K_{l_i} = H_i$.

We have that $(SH_*(K_s, J))$ is a filtered directed system.

Because every orbit is contained in a region of the form $l_{i-1} \leq r_N \leq l_i$ and

$$l_{i-1}(l_{i-1} + b) + b' - \frac{1}{2}l_i^2 \leq -\theta_N(X_H) - H \leq l_i(l_i + b') + b - \frac{1}{2}l_{i-1}^2$$

we have constants A, B so that any orbit starting on the level set $r_N = c$ must have action between Ac^2 and Bc^2 . Choose l_i so that $Al_i^2 \geq \lambda^2$. All orbits of H and K_{l_i} of action $\leq \lambda^2$ are contained inside the region $r_N \leq l_i$. The maximum principle (Corollary 9.2) ensures that all Floer trajectories connecting these orbits must be contained inside $r_N \leq l_i$. Hence $SH_*^{\leq \lambda^2}(H, J) = SH_*^{\leq \lambda^2}(K_{l_i}, J)$. We have a natural map:

$$SH_*^{\leq \lambda^2}(K_{l_i}, J) \rightarrow SH_*(K_{l_i}, J).$$

Composing these two maps gives us a natural map

$$p_{\lambda, i} : SH_*^{\lambda^2}(H, J) \rightarrow SH_*(K_{l_i}, J).$$

Because the filtered directed system maps for H are induced by inclusions and the filtered directed system maps for K_s are induced by non-decreasing families of Hamiltonians we have that $p_{\lambda, i}$ commutes with the filtered directed system maps as described in the statement of Lemma 4.19.

Now choose λ so that $Bl_i^2 \leq \lambda^2$. Then all orbits of K_{l_i} have action $\leq \lambda^2$. So the subcomplex of $CF_*^{\leq \lambda^2}(H, J)$ generated by orbits in the region $r_N \leq l_i$

is isomorphic to the chain complex $CF_*(K_{l_i}, J)$. Hence there is a natural morphism

$$p'_{i,\lambda} : SH_*(K_{l_i}, J) \rightarrow SH_*^{\leq \lambda^2}(H, J).$$

These morphisms also commute with the natural directed system maps. So by Lemma 4.19, $(SH_*^{\leq \lambda^2}(H, J))$ is isomorphic to $(SH_*(K_s, J))$ which in turn by Corollary 4.14 and Lemma 4.15 is isomorphic to $(SH_*(N, \lambda))$. $\square$

Proof. of Theorem 4.16. By Lemma 4.18, we get that $(SH_*^{\leq \lambda^2}(H, J))$ is isomorphic to $(SH_*^\#(T^*Q, \theta_Q, \lambda))$ for any quadratic admissible pair (H, J) . Using the metric on Q , we have a functional S defined on the loop space $\mathcal{L}Q$ given by sending a loop $l : S^1 = \mathbb{R}/\mathbb{Z} \rightarrow Q$ to $\sqrt{\int_0^1 |l'(t)|^2 dt}$. We have that $H_*(S^{-1}(-\infty, \lambda])$ is a filtered directed system where the directed system maps come from the natural inclusion maps. From [SW06, Corollary 1.2] we have a quadratic admissible pair (H, J) with the property that there is an isomorphism $SH_*^{\leq \lambda^2}(H, J, (\tau_Q, \omega_2)) \cong H_*(S^{-1}(-\infty, \lambda])$ for all λ . This isomorphism commutes with the directed system maps as well. Hence we get that $(SH_*^{\leq \lambda^2}(H, J, (\tau_Q, \omega_2)))$ is isomorphic to $(H_*(S^{-1}(-\infty, \lambda]))$ as directed systems. Let $L_{\text{par}}^{\leq \lambda}Q$ be the set of loops l of length $\leq \lambda$ such that $|l'(t)|$ is constant. This space is homotopic to $L^\lambda Q$. We have that $S(l)$ is equal to the length of l when $l \in L_{\text{par}}^{\leq \lambda}$. The result in [Ano81] tell us that the inclusion $L_{\text{par}}^{\leq \lambda}Q \hookrightarrow S^{-1}(-\infty, \lambda]$ is a homotopy equivalence. Hence $L^{\leq \lambda}$ is homotopy equivalent to $S^{-1}(-\infty, \lambda]$. We now get that $(SH_*^{\leq \lambda^2}(H, J, (\tau_Q, \omega_2)))$ is isomorphic to $(H_*(L^{\leq \lambda}Q))$ as directed systems. Hence $(SH_*^\#(T^*Q, \theta_Q, \lambda, (\tau_Q, \omega_2)))$ is isomorphic to $(H_*(L^\lambda))$ as directed systems. $\square$

4.2.1. The growth rate and fundamental group. Let G be a finitely generated group, and let $A := \{g_1, \dots, g_k\}$ be a set of generators for G . Let G^{cong} be the set of elements of G modulo conjugacy classes. Let r_i be the number of elements of G^{cong} which can be expressed as a product of at most i generators. We define $\Gamma^{\text{cong}}(G)$ to be $\overline{\lim}_i \frac{\log r_i}{\log i}$. This definition is independent of the choice of generators. The reason is as follows: Suppose $g'_1, \dots, g'_{k'}$ is another set of generators, then all we need to do is show that the growth rate associated to A is the same as the growth rate associated to the union of the generators: $B := \{g_1, \dots, g_k, g'_1, \dots, g'_{k'}\}$. Let r'_i be the number of elements of G^{cong} which can be expressed as a product of at most i elements of B . Then $r_i \leq r'_i$ because $A \subset B$. We have that there exists a $K \in \mathbb{N}$ such that each element of B can be expressed as a product of at most K elements of A . Therefore $r'_i \leq Kr_i$ for all i (where K is independent of i). Hence the growth rates are the same.

Lemma 4.20. *Let Q be a compact oriented manifold, then*

$$\Gamma(Q) \geq \Gamma^{\text{cong}}(\pi_1(Q)).$$

Proof. We need to show the following fact: there exists a constant P depending only on Q such that for every pair of free loops γ_1, γ_2 in Q , there exists a free loop γ and elements k_1 and k_2 of $\pi_1(Q)$ such that γ_i represents the conjugacy class $[k_i]$, and γ represents the conjugacy class $[k_1.k_2]$ with $l(\gamma) \leq l(\gamma_1) + l(\gamma_2) + P$. Here l denotes the length of a loop with respect to the metric g . This is done as follows: Choose a constant P such that for every pair of points $q_1, q_2 \in Q$, there exists a path joining them of length $\leq \frac{1}{2}P$. Also choose a basepoint $a \in Q$. Join γ_1 and γ_2 to a using paths of length $\leq \frac{1}{2}P$. This gives us elements $k_1, k_2 \in \pi_1(Q, a)$. The composition of such loops gives us a loop of length $\leq l(\gamma_1) + l(\gamma_2) + P$.

Let $g_1, \dots, g_k$ be generators for $\pi_1(Q, a)$. Let C be a constant greater than $l(g_i)$ for each i . For $C \geq 1$ we have that $\text{rank}(H_0^{\leq (C+P)c}(\mathcal{L}Q))$ is an upper bound for the number of loops of word length $\leq c$. Also the growth rate associated to $H_0^{\leq (C+P)c}(\mathcal{L}Q)$ is the same as the growth rate of $H_0^{\leq c}(\mathcal{L}Q)$. Hence $\Gamma(T^*Q, (\tau_Q, \omega_2))$ is an upper bound for $\Gamma^{\text{cong}}(\pi_1(Q))$. $\square$

Lemma 4.21. *Let G be the free product of 3 non-trivial groups then*

$$\Gamma^{\text{cong}}(G) = \infty.$$

Proof. of Lemma 4.21. Let $G = A \star B \star C$ where A, B, C are non-trivial. Let a, b, c be non-trivial elements in A, B, C . Choose a subset $I \subset \{1, \dots, k\}$. Let $q_I(i)$ be a function which is 1 if $i \in I$ and 0 otherwise. Let $a_I := bc \prod_{i=1}^k (a^{q_I(i)})(bc)^{1-q_I(i)}$.

Every element of G can be written uniquely in the form $f := \prod_{i=0}^l h_i$ where $h_i \in A$ or B or C and h_{i-1} is not in the same group A, B, C as h_i . There is an $l' < \frac{l}{2}$ such that $h_i = h_{l-i}^{-1}$ for all $i \leq l'$. The element $\prod_{i=l'+1}^{l-l'-1} h_i$ is called the *conjugation interior* of f . We can conjugate f by some element g so that the conjugation interior is of the form $\prod_{i=l'+1}^{l-l'-1} h_i$ where $h_{l'+1}$ is not in the same group A or B or C as $h_{l-l'-1}$. This is called a *standard conjugation interior*. If we have some element $\prod_{j=1}^{l'} h'_j$ as described above then a *rotation* of $\prod_{j=1}^{l'} h'_j$ is the operation where we replace it with $h'_l \prod_{j=1}^{l'-1} h'_j$ or we do the reverse.

If I conjugate f by any element g then the standard conjugation interior can only change by a sequence of rotations.

The conjugation interior of a_I is equal to a_I and it is standard. Hence if $a_{I'}$ is conjugate to a_I then $I = I' + j$ where we view $\{1, \dots, k\}$ as the cyclic group with k elements. We say that I' is a rotation of I if $I = I' + j$ for some $j \in \mathbb{Z}$. Also the word length of all these a_I 's with respect to the generators a, b, c are all the same. This word length is between $2 + k$ and $2 + 2k$. Hence the number of conjugacy classes of elements of G of word length between $2 + k$ and $2 + 2k$ in a, b, c is at least the number of subsets $I \subset \{1, \dots, k\}$ modulo rotation which is at least $\frac{2^k}{k}$. Hence the growth rate of G is bounded below by the growth rate of $\frac{2^k}{k}$ and hence is infinite. $\square$

5. COMPACTIFICATIONS OF ALGEBRAIC VARIETIES

5.1. Making the divisors orthogonal. Let (M, ω) be a symplectic manifold of dimension $2n$. Let $S_1, \dots, S_k$ be symplectic submanifolds of real co-dimension 2. For each $I \subset \{1, \dots, k\}$ we define S_I to be $\cap_{i \in I} S_i$. We say that $S_1, \dots, S_k$ are *symplectically intersecting* if they intersect transversally and S_I is a symplectic submanifold for each $I \subset \{1, \dots, k\}$. For any symplectic submanifold $S \subset M$, we define its ω -orthogonal bundle NS to be the vector subbundle of $TM|_S$ given by vectors u satisfying $\omega(u, v) = 0$ for each $v \in TS$.

Definition 5.1. We say that $S_1, \dots, S_k$ intersect positively if

- (1) They are symplectically intersecting.
- (2) Let $I \subset \{1, \dots, k\}$ be a disjoint union $I_1 \sqcup I_2$, N_1 the ω -orthogonal bundle of S_I inside S_{I_1} and N_2 the ω -orthogonal bundle of S_I inside S_{I_2} . The bundle $TS_I \oplus N_1 \oplus N_2$ is isomorphic to $TM|_S$. Each bundle TS_I, N_1 and N_2 has an orientation induced by ω and hence their direct sum does. We require that the natural orientation on this direct sum matches the orientation induced by ω^n on $TM|_S$.

Definition 5.2. Let $Q_1, \dots, Q_l$ be a collection of symplectic submanifolds of any dimension. Let U be any subset of M . We say that $Q_1, \dots, Q_k$ are orthogonal along U if for each $i, j \in \{1, \dots, k\}$ ($i \neq j$) and $x \in U \cap Q_i \cap Q_j$, the ω orthogonal normal bundle to Q_i at x is contained in TS_j . We just say they are orthogonal if they are orthogonal along M .

The aim of this section is to deform positively intersecting submanifolds so that they become orthogonal.

Lemma 5.3. Let $S_1, \dots, S_k$ be any finite set of positively intersecting symplectic submanifolds. There is a smooth family of positively intersecting symplectic submanifolds S_i^t such that

- (1) $S_i^0 = S_i$ for all $i \in \{1, \dots, k\}$.
- (2) All the S_i^1 intersect orthogonally.
- (3) $S_k^t = S_k$.

We need some preliminary lemmas and definitions before we prove the above lemma.

Lemma 5.4. Suppose we have k smooth families of symplectic submanifolds $S_1^t, \dots, S_k^t$ parameterized by $t \in [0, 1]$ and that $S_1^t, \dots, S_k^t$ are symplectically intersecting for each t . Suppose also that $S_1^0, \dots, S_k^0$ are positively intersecting. Then for each t , $S_1^t, \dots, S_k^t$ are also positively intersecting.

Proof. of Lemma 5.4. For $I \subset \{1, \dots, k\}$ let S_I^t be equal to $\cap_{i \in I} S_i^t$. If I is the disjoint union of I_1 and I_2 then we define N_i^t to be equal to be the ω -orthogonal bundle of S_I^t inside $S_{I_i}^t$. We have a smooth family of bundles $TS_I^t \oplus N_1^t \oplus N_2^t$, and the orientation of $TS_I^0 \oplus N_1^0 \oplus N_2^0$ agrees with that of

$TM|_{S_I^0}$. Hence because we have a smooth family of bundles, the orientation of $TS_I^t \oplus N_1^t \oplus N_2^t$ agrees with the orientation of $TM|_{S_I^t}$. $\square$

From now on, $\mathbb{R}^{2n}$ is the standard symplectic vector space with coordinates $x_1, y_1, \dots, x_n, y_n$ and symplectic form $\omega_{\text{std}} = \sum_{i=1}^n dx_i \wedge dy_i$.

Lemma 5.5. *Let $S_1, \dots, S_k$ be codimension 2 symplectic vector subspaces of $\mathbb{R}^{2n}$ such that $S_k = \{x_n, y_n = 0\}$ and where $S_1, \dots, S_k$ are positively intersecting. Then for all $\mu \geq 0$, $S_1, \dots, S_k$ are positively intersecting with respect to the new symplectic form $\omega_\mu := \omega_{\text{std}} + \mu dx_n \wedge dy_n$.*

Proof. of Lemma 5.5. Let $I \subset \{1, \dots, k\}$. If $k \in I$ then S_I is a subset of S_k , hence ω_μ restricted to S_I is equal to ω_{std} restricted to S_I and hence S_I is symplectic with respect to ω_μ for all $\mu \geq 0$. Now suppose that $k \notin I$. Then S_I is transverse to S_k . Let F be the ω_{std} orthogonal subspace of S_I to $S_I \cap S_k \subset S_I$. This is a two dimensional symplectic subspace. Let N be the ω_{std} orthogonal bundle to $S_I \cap S_k$ inside S_k . We know that $N \oplus (S_I \cap S_k) \oplus F$ has the same orientation as $\mathbb{R}^{2n}$ because $S_1, \dots, S_k$ are positively intersecting. Also $N \oplus (S_I \cap S_k)$ has the same orientation as S_k because N and S_I are ω_{std} orthogonal. Hence the orientation on F is the same as the orientation induced by restricting $dx_n \wedge dy_n$ to F . Hence ω_μ restricted to F is a volume form on F for all $\mu \geq 0$. Also F is still orthogonal to $S_I \cap S_k$ inside S_I with respect to ω_μ because $\omega_\mu(V, W) = \omega_{\text{std}}(V, W)$ for all vectors V, W where V is tangent to S_k . Hence ω_μ restricted to S_I is still a symplectic form.

So $S_1, \dots, S_k$ is symplectically intersecting with respect to ω_μ for all μ . Because $S_1, \dots, S_k$ are positively intersecting with respect to $\omega_0 = \omega_{\text{std}}$ we have by Lemma 5.3 that $S_1, \dots, S_k$ are all positively intersecting with respect to ω_μ for all $\mu \geq 0$. $\square$

Lemma 5.6. *Let $S_1, \dots, S_k$ be transversally intersecting codimension 2 vector subspaces of $\mathbb{R}^{2n}$ such that $S_k = \{x_n, y_n = 0\}$. Suppose that $S_1 \cap S_k, \dots, S_{k-1} \cap S_k$ are symplectically intersecting inside S_k , then for large enough $\mu \geq 0$ we have that $S_1, \dots, S_k$ are symplectically intersecting with respect to the symplectic form $\omega_\mu := \omega_{\text{std}} + \mu dx_n \wedge dy_n$.*

Proof. of Lemma 5.6. Let $I \subset \{1, \dots, k\}$. If $k \in I$ then S_I is a symplectic manifold with respect to ω_μ . From now on we will assume that $k \notin I$. We have that S_I intersects S_k transversally. Let $F \subset S_I$ be the vector subspace of S_I which consists of vectors which are ω_{std} orthogonal to $S_I \cap S_k$. First of all F is of dimension at least 2 because the set of vectors ω_{std} orthogonal to $S_I \cap S_k$ inside $\mathbb{R}^{2n}$ is $2(|I| + 1)$ dimensional and S_I is $2(n - |I|)$ dimensional. Here $|I|$ means the number of elements in I . Also F has dimension at most 2 because it must be orthogonal to $S_I \cap S_k$ inside S_I . Hence S_I is equal to $(S_I \cap S_k) \oplus F$ and ω_{std} restricted to S_I splits up under this direct sum as $\omega_1 \oplus \omega_2$. Because $S_I \cap S_k$ is symplectic by assumption we have that ω_1 is a symplectic form. Also $dx_n \wedge dy_n$ restricted to F is a non-degenerate 2-form because F is 2-dimensional and transverse to S_k . So for large enough μ ,

$\omega_2 + \mu dx_n \wedge dy_n|_F$ is a symplectic form on F . Hence $\omega_{\text{std}} + \mu dx_n \wedge dy_n$ is a symplectic form on S_I for large enough μ . Hence we have shown that S_I is symplectic for all I with respect to $\omega_{\text{std}} + \mu dx_n \wedge dy_n$ for large enough $\mu \geq 0$. $\square$

We have a parameterized version of this lemma that will be needed where we have a continuous family $S_1^q, \dots, S_{k-1}^q$ parameterized by $q \in Q$ where Q is some compact topological space. The result is that $S_1^q, \dots, S_{k-1}^q, S_k$ are symplectically intersecting with respect to ω_μ for μ sufficiently large.

Lemma 5.7. *Let S_k be a co-dimension 2 symplectic vector subspace of $\mathbb{R}^{2n}$. Let $A_1, \dots, A_{k-1}$ be symplectically intersecting symplectic vector subspaces of S_k . Let $\mathcal{B}$ be the space of $(k-1)$ -tuples of $2n-2$ dimensional symplectic vector subspaces $(S_1, \dots, S_{k-1})$ of $\mathbb{R}^{2n}$ such that $S_1, \dots, S_k$ are positively intersecting and such that $S_i \cap S_k = A_i$. If $\mathcal{B}$ is non-trivial then it deformation retracts onto the point $(A_1 + S_k^\perp, \dots, A_{k-1} + S_k^\perp)$ where $S_k^\perp$ is the symplectic orthogonal subspace to S_k .*

In particular we also get that $(A_1 + S_k^\perp, \dots, A_{k-1} + S_k^\perp)$ are also positively intersecting and hence $A_1, \dots, A_{k-1}$ are positively intersecting inside S_k .

Proof. of Lemma 5.7. By a linear symplectic change of coordinates we can assume that $S_k = \{x_n, y_n = 0\}$. We will show that $\mathcal{B}$ is weakly contractible and hence contractible. Let $S_1, \dots, S_{k-1}$ be a point in $\mathcal{B}$. The subspace $S_{\{1, \dots, k-1\}}$ is at least two dimensional and transverse to S_k . Let $W \subset S_{\{1, \dots, k-1\}}$ be a two dimensional symplectic vector subspace transverse to S_k . Such a subspace exists for the following reason: We have that $S_1, \dots, S_k$ intersect transversally and because these are codimension 2 vector subspaces we have that $k \leq n$. Hence $S_{\{1, \dots, k-1\}}$ has codimension at most $2(n-1)$ which means that it is at least 2 dimensional. This space is also transverse to S_k . Hence we can find such a two dimensional subspace W . The vector subspace W is also contained in S_i for each $1 \leq i \leq k-1$. Hence S_i is a direct sum of vector spaces $(S_i \cap S_k) \oplus W$ for each $1 \leq i \leq k-1$. Let $S_1^q, \dots, S_{k-1}^q$ be a family of points in $\mathcal{B}$ continuously parameterized by points q in some sphere S^m . Let $\phi^q : \mathbb{R}^2 \rightarrow \cap_{i=1}^{k-1} S_i^q$ be a continuous family of linear embeddings parameterized by q such that the image of ϕ^q is transverse to S_k . Now choose a continuous family of maps $\phi_t^q : \mathbb{R}^2 \rightarrow \mathbb{R}^{2n}$ parameterized by $(q, t) \in S^m \times [0, 1]$ such that they are linear embeddings transverse to S_k , $\phi_0^q = \phi^q$ and such that ϕ_1^q is a linear isomorphism to $S_k^\perp$. We now have a family of transversally intersecting (not necessarily symplectic) vector subspaces $S_i^{q,t} := A_i \oplus \text{image}(\phi_t^q)$. We have that $S_i^{q,0} = S_i^q$ and $S_i^{q,1} = A_i \oplus S_k^\perp$. By (a parameterized version of) Lemma 5.6, we have that $S_i^{q,t}$ is symplectically intersecting with respect to $\omega_\mu := \omega_{\text{std}} + \mu dx_n \wedge dy_n$ where $\mu \geq 0$ is very large (here we really need that $S_i^{q,t}$ is parameterized by a compact family and this is why we need to prove weak contractibility first). Let Φ_μ be a linear automorphism sending $(x_1, y_1, \dots, x_{n-1}, y_{n-1}, x_n, y_n)$ to

$(x_1, y_1, \dots, x_{n-1}, y_{n-1}, x_n, \frac{1}{1+\mu}y_n)$. Then $\Phi_\mu^* \omega_\mu = \omega_{\text{std}}$. This automorphism also preserves S_k and $S_k^\perp$, so $\Phi_\mu^* S_i^{q,1} = S_i^{q,1}$.

By Lemma 5.5 we have that $\Phi_{t\mu}^* S_i^{q,0}$ is positively intersecting for all $t \in [0, 1]$. We concatenate the isotopies $\Phi_{t\mu}^* S_i^{q,0}$ and $\Phi_\mu^* S_i^{q,t}$ giving us a new family of vector subspaces $(\sigma_1^{q,t}, \dots, \sigma_{k-1}^{q,t})$ defined as follows:

- (1) For $t \in [0, \frac{1}{2}]$, $\sigma_i^{q,t} = \Phi_{2t\mu}^* S_i^{q,0}$.
- (2) For $t \in [\frac{1}{2}, 1]$, $\sigma_i^{q,t} = \Phi_\mu^* S_i^{q,2t-1}$.

By Lemma 5.4 we get that $(\sigma_1^{q,t}, \dots, \sigma_{k-1}^{q,t}, S_k)$ are positively intersecting because $(\sigma_1^{q,0}, \dots, \sigma_{k-1}^{q,0}, S_k)$ are. Hence $(\sigma_1^{q,t}, \dots, \sigma_{k-1}^{q,t})$ are points in $\mathcal{B}$ parameterized by $q \in S^m$ starting at $S_1^q, \dots, S_{k-1}^q$ and ending at $A_1 \oplus S_k^\perp, \dots, A_k \oplus S_k^\perp$. Hence $\mathcal{B}$ is weakly contractible and hence contractible. $\square$

Lemma 5.8. *Let $\mathcal{S}_k$ be the space of k -tuples of positively intersecting $2n-2$ dimensional symplectic vector subspaces $S_1, \dots, S_k$ of $\mathbb{R}^{2n}$. Then $\mathcal{S}_k$ deformation retracts onto the space of positively intersecting ω_{std} orthogonal vector subspaces of $\mathbb{R}^{2n}$.*

Proof. of Lemma 5.8. Throughout the proof of the Lemma, we use the following fact: suppose that I have a fibration p whose fibers deformation retract to the fibers of a subfibration p' . Then the total space of p deformation retracts to the total space of p' .

We proceed by induction on dimension. Suppose this is true for all $\mathbb{R}^{2l}$ with $l < n$ and consider $\mathbb{R}^{2n}$. We define $\mathcal{S}_{k-1}^{S_k}$ as the space of positively intersecting $2n-4$ dimensional symplectic vector subspaces $A_1, \dots, A_{k-1}$ of S_k . Then this deformation retracts to the space of orthogonally intersecting subspaces by our induction hypothesis.

We have a fibration:

$$P : \mathcal{S}_k \twoheadrightarrow \mathcal{S}_1$$

sending $(S_1, \dots, S_k)$ to S_k . We have an inclusion $\mathcal{S}_{k-1}^{S_k} \hookrightarrow P^{-1}(S_k)$ sending $A_1, \dots, A_{k-1}$ to $(A_1 + S_k^\perp, \dots, A_{k-1} + S_k^\perp)$ where $S_k^\perp$ is the symplectic orthogonal subspace to S_k . So from now on we view $\mathcal{S}_{k-1}^{S_k}$ as a subspace of $P^{-1}(S_k)$ and the union $\cup_{S_k \in \mathcal{S}_1} \mathcal{S}_{k-1}^{S_k}$ as a subfibration of P . The fiber $P^{-1}(S_k)$ is also a fibration $Q : P^{-1}(S_k) \twoheadrightarrow \mathcal{S}_{k-1}^{S_k}$ where $Q(S_1, \dots, S_k) := (S_1 \cap S_k, \dots, S_{k-1} \cap S_k)$. The fiber of Q over a point $(A_1, \dots, A_{k-1})$ consists of subspaces $B_1, \dots, B_{k-1}$ positively intersecting S_k such that $B_i \cap S_k = A_i$. This deformation retracts to the point $(A_1 + S_k^\perp, \dots, A_{k-1} + S_k^\perp)$ by Lemma 5.7. Hence $P^{-1}(S_k)$ deformation retracts to the space $\mathcal{S}_{k-1}^{S_k}$ which by our induction hypothesis deformation retracts to the space of orthogonally intersecting subspaces of $\mathbb{R}^{2n-2}$. This implies that $\mathcal{S}_k$ deformation retracts to the space of orthogonally intersecting subspaces. This proves our Lemma. $\square$

Lemma 5.9. *Let $S \subset M$ be a symplectic submanifold of M . Suppose that the ω -orthogonal bundle $\pi_S : NS \twoheadrightarrow S$ has structure group $G \subset U(n)$. Then*

there is a neighbourhood US and a projection $p : US \rightarrow S$ whose fiber is symplectomorphic to the ball B_ϵ of radius ϵ and such that its structure group is G and whose fibers are orthogonal to S .

There is also a vector field L on US whose flow ϕ_t is well defined for all negative t and such that $e^t(\phi_{-t})^*\omega_{US} = \omega_{US} + (e^t - 1)\pi^*\omega_S$ where ω_{US} and ω_S are the symplectic forms on US and S respectively (here S is identified with the zero section). Inside the fiber $B_\epsilon \subset \mathbb{C}^n$, this vector field is equal to $\sum_i \frac{r_i}{2} \frac{\partial}{\partial r_i}$ where $(r_i, d\theta_i)$ are polar coordinates for the i th factor of $\mathbb{C}$ in $\mathbb{C}^n$.

Proof. of Lemma 5.9. This proof is very similar to the proof of [MS98, Theorem 6.3]. Let $(V_\alpha)_{\alpha \in A}$ be a finite covering of S with trivializations $V_\alpha \times \mathbb{C}^n$ of the symplectic fibration. Choose a partition of unity $\rho_\alpha : V_\alpha \rightarrow \mathbb{R}$ subordinate to this cover. Let $\pi_{\mathbb{C}^n}$ be the natural projection from $V_\alpha \times \mathbb{C}^n$ to $\mathbb{C}^n$. We define $\sigma_\alpha := \sum_i r_i^2 d\theta_i$ where (r_i, θ_i) are polar coordinates for the i th $\mathbb{C}$ factor of the fiber $\mathbb{C}^n$. We write $\tau := \sum_\alpha d((\rho_\alpha \circ \pi_S)\sigma_\alpha)$ which is equal to

$$\tau = \sum_\alpha d(\rho_\alpha \circ \pi_S) \wedge \sigma_\alpha + \rho_\alpha \circ \pi_S d\sigma_\alpha.$$

Because σ_α restricted to the zero section is zero and TS is in the kernel of $d\sigma_\alpha$, we have that $\omega_{US} := \pi_S^*\omega|_S + \tau$ is a symplectic form on a small neighbourhood of the zero section. This is because $\|d(\rho_\alpha \circ \pi_S) \wedge \sigma_\alpha\|$ is small relative to $\|\rho_\alpha \circ \pi_S d\sigma_\alpha\|$ and $\|\pi_S^*\omega|_S\|$ near the zero section. Let NS_ϵ be the open subset consisting of vectors of modulus less than ϵ where ϵ is small enough so that ω_{US} is still a symplectic form on NS_ϵ . We have by a Moser theorem [MS98, Theorem 3.3] that for some ϵ small enough, a neighbourhood of S is symplectomorphic to NS_ϵ . This has structure group G and fibers B_ϵ .

The fibers are orthogonal to S because if we have any vector v tangent to S and another one w tangent to the fiber, then $\pi_S^*\omega|_S(v, w) = 0$ because w is in the kernel of this symplectic form. Also $d(\rho_\alpha \circ \pi_S) \wedge \sigma_\alpha(v, w) = 0$ because σ_α vanishes on S . Finally $d\sigma_\alpha(v, w)$ vanishes because TS is in the kernel of $d\sigma_\alpha$. Hence $\omega_{US}(v, w) = 0$ and hence the fibers are orthogonal to S .

Let L be the vector field on NS_ϵ which is tangent to the fibers of π_S and equal to $\sum_i \frac{r_i}{2} \frac{\partial}{\partial r_i}$ on the fibers. This is well defined because $\sum_i \frac{r_i}{2} \frac{\partial}{\partial r_i}$ is invariant under the $U(n)$ action on $\mathbb{C}^n$. We have that the Lie derivative of $\rho_\alpha \circ \pi_S \cdot \sigma_\alpha$ with respect to L is $\rho_\alpha \circ \pi_S \cdot \sigma_\alpha$ (because the flow of L does not change $\rho_\alpha \circ \pi_S$). Hence $(\phi_t)^*\tau = e^t\tau$ where ϕ_t is the flow of L . Hence

$$e^t(\phi_{-t})^*\omega_{US} = (e^t - 1)\omega|_S + \omega_{US}.$$

□

Lemma 5.10. *Let $p : US \rightarrow S$ be a fibration described in Lemma 5.9 and let $N \subset N'$ be open subsets of the zero section such that the closure of N is contained in N' (S here is compact and it may or may not have a boundary).*

Let $S_1, \dots, S_k$ be positively intersecting co-dimension 2 submanifolds such that

- (1) $\cap_i S_i$ is the zero section.
- (2) S_i intersects the fibers transversely.
- (3) S_i intersect orthogonally on $p^{-1}(N')$
- (4) $S_i \cap p^{-1}(q)$ is a symplectic vector subspace of $B_\epsilon \subset \mathbb{C}^n$ for each $q \in S$.

Then there is a family of manifolds S_i^t ($t \in [0, 1]$)

- (1) S_i^t are positively intersecting symplectic manifolds with respect to the symplectic form $\omega_{US} + Tp^*\omega_S$ for sufficiently large $T > 0$.
- (2) $\cap_i S_i^t$ is the zero section.
- (3) $S_i^0 = S_i$.
- (4) $S_i^t \cap p^{-1}(N) = S_i \cap p^{-1}(N)$.
- (5) S_i^1 are all orthogonal along the zero section.
- (6) $S_i^t = S_i$ outside some neighbourhood of the zero section.

Proof. of Lemma 5.10.

By Lemma 5.8, there is a family of submanifolds W_i^t of US transverse to all the fibers such that

- (1) $W_i^t \cap p^{-1}(q)$ are positively intersecting symplectic vector subspaces of $p^{-1}(q)$ for all $q \in S$.
- (2) W_i^t intersect each fiber of p transversely.
- (3) $W_i^0 = S_i$.
- (4) $W_i^t = S_i$ on $p^{-1}(N)$
- (5) $W_i^1 \cap p^{-1}(q)$ are orthogonal inside $p^{-1}(q)$ for all $q \in S_I$.

Note that W_i^t become positively intersecting symplectic submanifolds with respect to the symplectic form $\omega_{US} + Tp^*\omega_S$ for all sufficiently large $T > 0$.

For each $s \in [0, 1]$, we can construct (using a bump function) a smooth family of co-dimension 2 submanifolds $U_i^{s,t}$ ($t \in [0, 1]$) such that $U_i^{s,t} = W_i^s$ outside some closed subset of US containing the zero section, $U_i^{s,t} = W_i^t$ near the zero section and $U_i^{s,s} = W_i^s$. We can also assume that $U_i^{s,t} = W_i^s$ on $p^{-1}(N)$ because $W_i^s = W_i^t$ in this region. For each s , there is a constant $\delta_s > 0$ such that for all $t \in [s - \delta_s, s + \delta_s]$, $U_1^{s,t}, \dots, U_k^{s,t}$ are positively intersecting symplectic submanifolds with respect to $\omega_{US} + Tp^*\omega_S$ for all sufficiently large $T > 0$. We also assume that $U_i^{s,t}$ are transverse to all the fibers of p for all $t \in [s - \delta_s, s + \delta_s]$.

Let L be the vector field described in the statement of Lemma 5.9 and let ϕ_t be its flow. For any $S \geq 0$ we have that $\phi_{-S}(U_i^{s,t})$ is still a symplectic submanifold with respect to the symplectic form $\omega_{US} + Tp^*\omega_S$ because

$$e^S(\phi_{-S})^*(\omega_{US} + Tp^*\omega_S) = \omega_{US} + (T + 2e^S - 1)p^*\omega_S.$$

From now on we define $\phi_{-T}(U_i^{s,t})$ to be equal to $\phi_{-T}(U_i^{s,t})$ inside $\phi_{-T}(US)$ and equal to $U_i^{s,t}$ outside $\phi_{-T}(US)$. This is a well defined symplectic manifold because $U_i^{s,t} \cap p^{-1}(q)$ is linear outside some closed subset of US containing the zero section for all $q \in S$ and the flow of L is tangent to any

submanifold with this property. We also have that $\phi_{-T}(U_i^{s,t})$ are all positively intersecting symplectic submanifolds. Also because L is tangent to the fibers of p , we have that $\phi_{-T}(U_i^{s,t})$ is transverse to all the fibers.

By compactness, there exists a finite sequence $s_1 < s_2 < \dots < s_l$ such that the intervals $(s_i, s_i + \delta_{s_i})$ cover $(0, 1]$. This is because we can choose $s_1 = 0$ and then cover $[\delta_{s_1}, 1]$ with the open intervals $(s_i, s_i + \delta_{s_i})$ by compactness. We assume that $s_l = 1$ and $s_{i+1} \in (s_i, s_i + \delta_{s_i})$.

The submanifold $\phi_{-T}(U_i^{s,t})$ is still equal to W_i^s outside some closed subset of US containing the zero section and equal to W_i^t near the zero section. Suppose we have found a family S_i^t ($t \in [0, s_{m-1}]$) such that $S_i^t = S_i$ outside some relatively compact open set O and such that $S_i^{s_{m-1}} = W_i^{s_{m-1}}$ near 0. We also suppose that $S_i^t = S_i$ on $p^{-1}(N)$. We now consider s_m . Let $O_m \subset O$ be an open set containing the zero section such that $S_i^{s_{m-1}} = W_i^{s_{m-1}}$ on a neighbourhood of the closure of O_m for all i . For T large enough, we have $G_t := \phi_{-T}(U_i^{s_{m-1},t})$ is equal to $W_i^{s_{m-1}}$ inside a neighbourhood of the closure of $US \setminus O_m$. Note that G_t is well defined for $t \in [s_{m-1}, s_m]$ because $(s_{m-1} - \delta_{s_{m-1}}, s_{m-1} + \delta_{s_{m-1}})$ contains $[s_{m-1}, s_m]$. We define S_i^t for $t \in [s_{m-1}, s_m]$ to be equal to $S_i^{s_{m-1}}$ outside O_m and equal to G_t inside O_m . This is a manifold because these manifolds agree on the boundary of O_m and just outside O_m . Also we have that $S_i^{s_m}$ is equal to $W_i^{s_m}$ near 0 because $U_i^{s_{m-1},s_m}$ does. We have that $S_i^t = S_i$ on $p^{-1}(N)$ because $U_i^{s,t}$ has this property and the flowing along L preserves this property.

Hence by induction we have constructed a family S_i^t equal to S_i outside O and such that $S_i^{s_l} = S_i^1$ is equal to W_i^1 near the zero section. Because W_i^1 are orthogonal at the zero section, we get that S_i^1 are also orthogonal at the zero section. Finally $S_i^t = S_i$ on $p^{-1}(N)$. This proves the Lemma. $\square$

Lemma 5.11. *Let $S_1, \dots, S_k$ be positively intersecting symplectic manifolds in M and $S'_1, \dots, S'_k$ in M' . Let NS_i, NS'_i be their respective symplectic normal bundles. Suppose that there is a series of symplectomorphisms $\phi_i : S_i \rightarrow S'_i$ and symplectic bundle isomorphisms $N\phi_i : NS_i \rightarrow NS'_i$ covering ϕ_i such that ϕ_i, ϕ_j and $N\phi_i, N\phi_j$ agree when restricted to $S_i \cap S_j$ for all i, j .*

Then there is a symplectomorphism Φ from a neighbourhood of $\cup_i S_i$ to a neighbourhood of $\cup_i S'_i$ such that $\Phi|_{S_i} = \phi_i$ and $D\Phi|_{NS_i} = N\phi_i$.

Suppose in addition that a neighbourhood NS_I of S_I is identical to a neighbourhood of NS'_I and that for $i \in I$, ϕ_i and $N\phi_i$ restricted to S_I is the identity map for all i . Then we may assume that Φ is Hamiltonian near S_I and C^1 small.

Proof. of Lemma 5.11. We will first construct a diffeomorphism Ψ from a neighbourhood of $\cup_i S_i$ to a neighbourhood of $\cup_i S'_i$ such that $\Psi|_{S_i} = \phi_i$ and $D\Psi|_{NS_i} = N\phi_i$ (i.e. every vector v in $N_x S_i$ is sent to a vector in $N_{\phi_i(x)} S'_j$ equal to $N\phi_i(v)$). We do this by induction: Suppose that we have found such a symplectomorphism Ψ' on a neighbourhood of $\cup_{i=1}^{m-1} S_i$ and consider $\cup_{i=1}^m S_i$. By using an exponential map we have a diffeomorphism

$\tilde{\phi}_m$ from a neighbourhood of S_m to a neighbourhood of S'_m such that $D\tilde{\phi}_m$ induces the morphism $N\phi_m$. On $\cup_{i=1}^{m-1}(S_i \cap S_m)$ we have that $\tilde{\phi}_m$ and Ψ' agree. Also $D\tilde{\phi}_m|_{\cup_{i=1}^{m-1} \cap S_m}$ is equal to $D\Psi'|_{\cup_{i=1}^{m-1} \cap S_m}$. Hence if we look at their graphs inside $M \times M'$, they are tangent along $(\cup_{i=1}^{m-1}(S_i \cap S_m) \times M'$ so we can perturb the graph of $\tilde{\phi}_m$ by a C^1 small amount so that it still stays the graph of a diffeomorphism ϕ' and such that it agrees with Ψ' on a neighbourhood of $\cup_{i=1}^{m-1}(S_i \cap S_m)$. Hence we can extend Ψ' over S_m with the appropriate properties.

Finally, we can use a standard Moser deformation argument to deform Ψ into a symplectomorphism Φ such that the differential of Φ at $\cup_i S_i$ agrees with the differential of Ψ at $\cup_i S_i$. This ensures that Φ has the properties we want.

Now suppose in addition that a neighbourhood NS_I of S_I is identical to a neighbourhood of NS'_I and that for $i \in I$, ϕ_i and $N\phi_i$ restricted to S_I is the identity map for all i . We can construct a smooth family S_i^t $t \in [0, 1]$ of positively intersecting submanifolds which are C^1 close to each other so that on a small neighbourhood of S_I , S_i^0 is equal to S_i for all i . Also we want that globally, S_i^1 is equal to S'_i for all i . We can also assume that they are all symplectomorphic to each other, so we have a smooth family of symplectomorphisms ϕ_i^t from S_i to S_i^t and also a smooth family of normal bundle maps $N\phi_i^t$ so that when $t = 1$ they all coincide with ϕ_i and $N\phi_i$. By a parameterized version of the above discussion we have a smooth family of symplectomorphisms Φ^t from a neighbourhood of $\cup_i S_i$ to a neighbourhood of $\cup_i S_i^t$ such that $\Phi^t|_{S_i} = \phi_i^t$ and $D\Phi^t|_{NS_i} = N\phi_i^t$. In addition we can assume that Φ^0 is the identity map near S_I (because we can choose our associated diffeomorphism Ψ to have this property). Because ϕ_i and $N\phi_i$ are the identity map on S_I we have that Φ^1 is C^1 small near S_I . Let $V_t := \frac{d}{dt}\Phi^t$ be a smooth family of symplectic vector fields defined near S_I . These are Hamiltonian vector fields near S_I because $V_t = 0$ on S_I and a small neighbourhood of S_I deformation retracts onto S_I . Hence Φ^1 is a Hamiltonian symplectomorphism near S_I because it is Hamiltonian isotopic to Φ^0 near S_I which is the identity map. Hence $\Phi := \Phi^1$ has the required properties. $\square$

Lemma 5.12. *Let (M, ω) be a symplectic manifold and let S be a compact symplectic submanifold and N, N' open subsets of M such that the closure of N is contained in N' . Let $S_1, \dots, S_k$ be positively intersecting symplectic submanifolds with $\cap_i S_i = S$ and such that they are orthogonal along S , and such that they are also orthogonal in N' . Then there exists a C^1 small perturbation of $S_1, \dots, S_k$ to $S'_1, \dots, S'_k$ such that $S'_1, \dots, S'_k$ are orthogonal on a small neighbourhood of S , and $S'_1 = S_1, \dots, S'_k = S_k$ outside a small neighbourhood of S and also inside N . We may also arrange that $TS'_i = TS_i$ along S .*

Proof. of Lemma 5.12. We prove this by induction on the dimension of M . So we suppose it is true in lower dimensions. Choose a small neighbourhood US of S as in Lemma 5.9 where $p : US \rightarrow S$ is a symplectic fibration whose fibers are orthogonal to S .

Consider the positively intersecting submanifolds $(S_i \cap S_k)_{i=1, \dots, k-1}$ inside S_k . By our induction hypothesis we can perturb them by a C^1 small amount to $V_1, \dots, V_{k-1}$ such that:

- (1) $V_i = S_i$ inside $N'' \subset S_k$ and also outside a small neighbourhood $M \subset S_k$ of $S \cap S_k$. Here N'' is an open subset whose closure is contained in $N' \cap S$ and such that it contains closure of $N \cap S_k$.
- (2) V_i intersect orthogonally along a small neighbourhood of $S \cap S_k$.

Let q be a projection from a small neighbourhood of S_k to S_k whose fibers are orthogonal to S_k . Let $W_i := q^{-1}(V_i)$ for $i = 1, \dots, k-1$. These symplectic submanifolds are tangent to S_i along S and orthogonal on a neighbourhood $p^{-1}(S)$ of S . Because the normal bundle to S_k is two dimensional and S_i, W_i are orthogonal to S_k in the region $(S_k \cap N'')$ and of co-dimension 2, we have that their tangent spaces coincide in this region. By Lemma 5.11 there is a C^1 small Hamiltonian symplectomorphism (defined near S_k) sending W_i to S_i for each i in the region $p^{-1}(N'')$ and fixing S_k . Choose a neighbourhood O of S small enough so that $N \cap O \subset p^{-1}(S_k \cap N'')$. This means that $W_i = S_i$ inside $N \cap O$. Hence we can perturb S_i for $i = 1, \dots, k-1$ by a C^1 small amount to S'_i so that it coincides with W_i on a small neighbourhood of S (inside O) and is equal to S_i outside O and is unchanged in N . This means that $S'_i = S_i$ inside N and S'_i is orthogonal on a neighbourhood of S . Also $TS'_i = TS_i$ along S . Hence S'_i has the properties we want. $\square$

Lemma 5.13. . *Let $S_1, \dots, S_k$ be positively intersecting symplectic submanifolds inside some symplectic manifold (M, ω) and let $I \subset \{1, \dots, k\}$. Suppose that $S_1, \dots, S_k$ intersect orthogonally on some neighbourhood N of $S_I \cap (\cup_{i \notin I} S_i)$.*

Then there is a family of positively intersecting symplectic manifolds S_i^t ($t \in [0, 1]$) with

- (1) $S_i^0 = S_i$.
- (2) $S_i^t = S_i$ for $i \notin I$.
- (3) $S_i^t = S_i$ on some open subset containing $S_I \cap (\cup_{i \notin I} S_i)$.
- (4) S_i^1 are all orthogonal on an arbitrarily small neighbourhood NS_I of S_I .
- (5) $S_i^t = S_i$ outside an arbitrarily small neighbourhood of the closure of NS_I .
- (6) $\cap_{i \in I} S_i^t = S_I$.

Proof. of Lemma 5.13. By Lemma 5.9, there is a neighbourhood US of S_I and a projection $p_I : US \rightarrow S_I$ such that the fibers of p_I are symplectomorphic to B_ϵ and whose structure group is $U(n)$. There is also a vector field

L tangent to the fibers of p_I such that $e^t(\phi_{-t})^*\omega = \omega + (e^t - 1)p^*\omega|_S$ and such that it radial in the fibers. We define $S := S_I$ and $\omega_S := \omega|_S$.

We prove the Lemma now by reducing it to the linear case in Lemma 5.10. For each $i \in I$, consider the following manifold:

$$T_i := \cup_{q \in S} T_0(S_i \cap p^{-1}(q)).$$

Here $T_0(S_i \cap p^{-1}(q))$ means the tangent space at 0 of $S_i \cap p^{-1}(q)$ inside the tangent space of the linear fiber $p^{-1}(q)$ at 0 which is canonically identified with $p^{-1}(q)$ (because $p^{-1}(q)$ is an open ball in $\mathbb{C}^n$). We have that T_i is symplectic near S_I . By a parameterized version of Lemma 5.11 there is a C^1 small diffeomorphism preserving the fibers of p_I sending T_i to S_i in the region $p^{-1}(N)$ for all $i \in I$ (possibly after shrinking N slightly) and such that it is a symplectomorphism when restricted to each fiber. We push forward the $U(n - |I|)$ structure group of $p_I : US \rightarrow S_I$ via this fiberwise diffeomorphism as well so that S_i restricted to each fiber $p^{-1}(q)$ is linear for each $q \in N$.

For each $i \in I$, we can perturb S_i by a C^1 small amount (without moving S_I) so that $S_i = T_i$ on a small neighbourhood P of the zero section and so that the S_i are all still positively intersecting symplectic submanifolds. We can also assume that this perturbation only happens outside some neighbourhood of $S_I \cap (\cup_{i \notin I} S_i)$.

So from now on (after shrinking US) we can assume that $S_i \cap p^{-1}(q)$ is linear inside $p^{-1}(q)$ for all q and $i \in I$. This means we have a codimension 0 submanifold $\bar{S} \subset S_I$ with boundary disjoint from $S_I \cap (\cup_{i \notin I} S_i)$ such that $(S_i)_{i \in I}$ are orthogonal away from $\bar{S}$ and on a neighbourhood $\mathcal{N}$ of the boundary of $\bar{S}$. Let $U\bar{S}$ be equal to $p^{-1}(\bar{S})$.

By Lemma 5.10 (using $U\bar{S}$) we can find a family of submanifolds S_i^t so that

- (1) S_i^t are positively intersecting symplectic manifolds with respect to the symplectic form $\omega_{US} + Tp^*\omega_S$ for sufficiently large $T > 0$.
- (2) $\cap_i S_i^t$ is the zero section.
- (3) $S_i^0 = S_i$.
- (4) $S_i^t \cap p^{-1}(\mathcal{N}) = S_i \cap p^{-1}(\mathcal{N})$.
- (5) S_i^1 are all orthogonal along the zero section.
- (6) $S_i^t = S_i$ outside some small neighbourhood of S .

We define $\phi_{-T}(S_i^t)$ to be equal to $\phi_{-T}(S_i^t)$ inside $\phi^{-T}(US)$ and S_i^t outside $\phi^{-T}(US)$. These are also positively intersecting submanifolds with the properties stated above (because $\phi_{-T}^*\omega_{US} = \omega_{US} + (1 - e^{-T})p^*\omega|_S$).

The problem is we want these symplectic manifolds to be orthogonal on a neighbourhood of S_I . But this can be done by perturbing $\phi_{-T}(S_i^t)$ by a C^1 small amount inside $U\bar{S}$ (by Lemma 5.12), hence $\phi_{-T}(S_i^t)$ has all the properties we want. $\square$

In the previous Lemma we have that $\cap_{i \in I} S_i^t = S_I$. We also have a smooth family of diffeomorphisms Ψ_t from S_i to S_i^t which are the identity on S_I and

outside a small neighbourhood of S_I . Because a neighbourhood of S_I inside S_i^t deformation retracts to S_I , we have that any 2-cycle in S_i^t near S_I is homologous to a 2-cycle in S_I . This means that the integral of $\omega|_{S_i}$ over this 2-cycle C is the same as the integral of $\omega|_{S_i^t}$ over $\Psi_*(C)$. Also away from this neighbourhood we have that $S_i^t = S_i$ so any 2-cycle evaluated on $\omega|_{S_i}$ is the same as the one evaluated on $\Psi^*\omega_{S_i^t}$. All of this implies that $\Psi_t^*([\omega]|_{S_i^t})$ is equal to $[\omega]|_{S_i}$ and by a Moser theorem we can then ensure that this is a symplectomorphism if the S_i are compact. Hence (by using another Moser theorem) there is a smooth family of symplectomorphisms $P_i^t : \text{Nhd}(S_i) \rightarrow \text{Nhd}(S_i^t)$ where Nhd means ‘a small neighbourhood of’. These symplectomorphisms fix S_I and hence by a similar cohomological argument (as explained earlier), we have that P_i^t is in fact a Hamiltonian symplectomorphism, so by using a cutoff function we can extend these symplectomorphisms to Hamiltonian symplectomorphisms: $P_i^t : M \rightarrow M$.

Proof. of Lemma 5.3. We basically proceed by induction on subsets I where $I \subset \{1, \dots, k\}$. In order to do induction, we need a total order on this finite set. Here is the following total order: We say that $I \prec J$ when

- (1) $|I| > |J|$ or
- (2) $|I| = |J|$ with $I \neq J$ and the highest number in $J \setminus (J \cap I)$ is smaller than the highest number in $I \setminus (J \cap I)$.

Fix some $I \subset \{1, \dots, k\}$. Suppose by our induction hypothesis, we have deformed $S_1, \dots, S_k$ through positively intersecting symplectic manifolds so that they are orthogonal on a small neighbourhood N of $\cup_{J \prec I} S_J$ and consider S_I . By Lemma 5.13, we can deform S_i through positively intersecting submanifolds S_i^t such that:

- (1) $S_i^0 = S_i$.
- (2) $S_i^t = S_i$ for $i \notin I$.
- (3) S_i^1 are all orthogonal on some small neighbourhood of S_I .
- (4) $S_i^t = S_i$ outside an arbitrarily small neighbourhood of S_I .

Because $S_i^t = S_i$ outside an arbitrarily small neighbourhood O of S_I , we can assume that $(O \cap S_j) \setminus N$ is empty for all $j \notin I$. This means that S_i^1 are still orthogonal along N and also on some small neighbourhood of S_I .

Hence by induction we have proven that we can deform S_i through positively intersecting symplectic submanifolds so that they are orthogonal on a neighbourhood of $\cup_i S_i$ and hence are orthogonal everywhere.

Let $(S'_i)^t$ be this deformation. The problem is that $(S'_k)^t$ is not equal to S_k for all t . The paragraph before this proof tells us that there is a sequence of symplectomorphisms P_k^t sending S_k to $(S'_k)^t$ such that P_k^0 is the identity map. So we can pull back $(S'_i)^t$ via P_k^t for all i and this ensures that we get a family of positively intersecting submanifolds S_i^t such that S_i^t are all positively intersecting, $S_i^0 = S_i$ and $S_k^t = S_k$. $\square$

5.2. Making the smooth affine variety nice at infinity.

Lemma 5.14. *Suppose that $S_1, \dots, S_k$ are positively intersecting codimension 2 symplectic submanifolds of (M, ω) such that they are also orthogonal. There exist small neighborhoods US_i of S_i and projections $\pi_i : US_i \rightarrow S_i$ such that*

- (1) *For $1 \leq i_1 < i_2 < \dots < i_l \leq k$,*

$$\pi_{i_l} \circ \dots \circ \pi_{i_1} : \cap_{j=1}^l US_{i_j} \rightarrow S_{\{i_1, \dots, i_l\}}$$

has fibers that are symplectomorphic to $\Pi_{j=1}^l \mathbb{D}_\epsilon$ where $\mathbb{D}_\epsilon$ is the disk of radius ϵ .

- (2) *If we look at a fiber $\Pi_{j=1}^l \mathbb{D}_\epsilon$ of $\pi_{i_l} \circ \dots \circ \pi_{i_1}$, then for $1 \leq m \leq l$, π_{i_m} maps this fiber to itself. It is equal to the natural projection*

$$\Pi_{j=1}^l \mathbb{D}_\epsilon \rightarrow \Pi_{j=1, j \neq m}^l \mathbb{D}_\epsilon$$

eliminating the m th disk $\mathbb{D}_\epsilon$.

- (3) *The symplectic structure on US_i induces a natural connection for $\pi_{i_l} \circ \dots \circ \pi_{i_1}$ given by the ω orthogonal vector bundles to the fibers. We may require the associated parallel transport maps to be elements of $U(1) \times \dots \times U(1)$ where $U(1)$ acts on the disk $\mathbb{D}_\epsilon$ by rotation.*

Proof. of Lemma 5.14. Suppose we have a bundle $p : V \rightarrow B$ with a $\prod_{i=1}^m U(1)$ structure where the fiber is $\mathbb{D}_\epsilon^m$ where $\mathbb{D}_\epsilon$ is the ϵ ball in $\mathbb{C}$ and the m 'th copy of $U(1)$ rotates the m 'th $\mathbb{D}_\epsilon$ factor in $\mathbb{D}_\epsilon^m$. Suppose that the base B has a symplectic structure ω_B . We can construct a symplectic structure on the total space V as follows: Let $V_i \subset V$ be the subbundle whose fiber is the subset is the i 'th copy of $\mathbb{D}_\epsilon$ in $\mathbb{D}_\epsilon^m$. This has a $U(1)$ structure group. By Lemma 5.9, there is a symplectic structure ω_{V_i} on V_i such that the fibers of V_i have the standard symplectic structure on $\mathbb{D}_\epsilon \subset \mathbb{C}$. We can ensure that the parallel transport maps are in $U(1)$ as well as follows: On V_i , there is an S^1 action $A : S^1 \rightarrow \text{diffeo}(V_i)$ such that it fixes the map p and rotates the fibers of V_i (i.e. it corresponds to the action given by rotating the fiber $\mathbb{D}_\epsilon \subset \mathbb{C}$). We define

$$\bar{\omega}_{V_i} := \int_{S^1} A(t)^* \omega dt.$$

This is a symplectic form if we shrink $\epsilon > 0$ a bit. Also the new symplectic form on V_i ensures that the parallel transport maps are in $U(1)$. Let $P_i : V \rightarrow V_i$ be the natural projection to V_i where $(a_1, \dots, a_m)$ in the fiber $\mathbb{D}_\epsilon^m$ is projected to the i 'th $\mathbb{D}_\epsilon$ factor. We define $\omega_V := \frac{1}{m} \sum_{i=1}^m P_i^* \bar{\omega}_{V_i}$. Any symplectic structure on a bundle with a $\prod_{i=1}^m U(1)$ structure group as described above is said to be *bundle compatible*. The good thing about bundle compatible symplectic structures is that the maps P_i satisfy all the properties as described in the start of this Lemma (where π_i is replaced by P_i). Also if we have an open subset U of B and bundle compatible symplectic structures on $p^{-1}(U)$ then by using similar methods and partitions of unity, we can ensure that there is a bundle compatible symplectic form on V

which coincides with the symplectic form on $p^{-1}(U)$ possibly after shrinking $p^{-1}(U)$ a tiny bit. We will call the maps P_i the *divisor projections*.

In order to prove our Lemma, we basically proceed by induction on subsets I where $I \subset \{1, \dots, k\}$. In order to do induction, we need a total order on this finite set. Here is the following total order: We say that $I \prec J$ when

- (1) $|I| > |J|$ or
- (2) $|I| = |J|$ with $I \neq J$ and the highest number in $J \setminus (J \cap I)$ is smaller than the highest number in $I \setminus (J \cap I)$ or

Let I_L be the L th subset in this total order. If $I \subset \{1, \dots, k\}$ then we write p_I to mean the map

$$\pi_{i_l} \circ \dots \circ \pi_{i_1} : \cap_{j=1}^l US_{i_j} \rightarrow S_{\{i_1, \dots, i_l\}}$$

where $I = \{i_1, \dots, i_l\}$ and $i_1, \dots, i_l$ are distinct. Suppose by our induction hypothesis there exists a neighbourhood U_{L-1} of $\cup_{i \leq L-1} S_{I_i}$ such that for each i with $1 \leq i \leq k$, we have a neighbourhood $U_{L-1} \cap US_i$ of S_i and a projection map $\pi_i : U_{L-1} \cap US_i \rightarrow S_i \cap U_{L-1}$ such that for each $I \subset \{1, \dots, k\}$, the map p_I has the structure of a $\prod_{i \in I} U(1)$ bundle which is bundle compatible with the symplectic structure. We will also assume that the maps π_i are the associated divisor projections for p_I locally around S_I inside U_{L-1} . Even though US_i has not been constructed yet, we will use the notation $U_{L-1} \cap US_i$ for the part of US_i that has been constructed inside U_{L-1} . We write $I := I_L$. We want to extend U_{L-1} to U_L containing S_I and $US_i \cap U_{L-1}$ to $US_i \cap U_L$ along with the maps π_i so that they satisfy the properties as stated above. The normal bundle of S_I has a natural $\prod_{i \in I} U(1)$ structure group because its tangent bundle is the intersection $\cap_i TS_i$. Hence by the previous discussion and using an exponential map, we can extend the maps π_i over a neighbourhood and we can extend the symplectic structure $\omega|_{U_{L-1}}$ to ω' over some neighbourhood of S_I so that:

- (1) π_I has the structure of a $\prod_{i \in I} U(1)$ bundle with fiber $\prod_{i \in I} \mathbb{D}_\epsilon$.
- (2) The symplectic structure ω' is bundle compatible.
- (3) the maps π_i are the associated bundle projections.

The symplectic structure ω' coincides with the symplectic structure ω_M on TS_I , and the symplectic normal bundles are the same although the symplectic form on one normal bundle is a positive scalar multiple of the other. We can make the symplectic forms coincide exactly along S_I by pushing forward ω' by a diffeomorphism induced by a vector field which is 0 on S_I and which is tangent to the normal bundle so that it rescales the symplectic form ω' on this normal bundle so it coincides with ω_M . Hence by a Moser theorem we can ensure that ω' coincides with ω_M (we have to deform the maps π_i as well). The problem is that because we have deformed the maps π_i , the image of π_i might not coincide with S_i away from $\cup_{j \notin I} S_j$ anymore. But by Lemma 5.11, there is a symplectomorphism which is the identity near $\cup_{j \notin I} S_j$ moving all the π_i 's so that their image is in S_i . Hence π_i has all the properties we want and is defined on a neighbourhood U_L of S_I .

Hence by induction we have maps π_i defined on some neighbourhood of $\cup_i S_i$ with the properties stated as in the Lemma. $\square$

Let $S_1^t, \dots, S_k^t$ ($t \in [0, 1]$) be a smooth family of codimension 2 symplectic submanifolds such that for each fixed t , $S_1^t, \dots, S_k^t$ are positively intersecting. We write $S^t := \cup_{i,j} (S_i^t \cap S_j^t)$.

Lemma 5.15. *There is a smooth family of symplectomorphisms Φ_t from $M \setminus S^0$ to $M \setminus S^t$ sending $S_i^0 \setminus S^0$ to $S_i^t \setminus S^t$ for each i .*

In particular this means that there is a smooth family of symplectomorphisms induced by Φ_t from $M \setminus (\cup_i S_i^0)$ to $M \setminus (\cup_i S_i^t)$.

We first need a preliminary Lemma:

Lemma 5.16. *Suppose M is any symplectic manifold (open or closed without boundary) and let $\iota_t : S \hookrightarrow M$ ($t \in [0, 1]$) be any smooth family of proper symplectic embeddings of the symplectic manifold S (without boundary). Then there is a family of Hamiltonian symplectomorphisms $\phi_t : M \rightarrow M$ such that ϕ_0 is the identity and ϕ_t sends the image of ι_0 to the image of ι_1 .*

Proof. of Lemma 5.16. By a Moser theorem, there is a neighbourhood U of $\iota_0(S)$ (which gets very thin near infinity) and a smooth family of symplectic embeddings $\iota'_t : U \hookrightarrow M$ such that $\iota'_t|_S = \iota_t$. We can also ensure that there is a smooth submersion $\pi : U \rightarrow S$ whose fibers are all ω_M -orthogonal to S . We will also assume that U deformation retracts to S .

For each t , there is a symplectic vector field V_t on $\iota'_t(U)$ defined as $\frac{d}{dt}(\iota'_t(p))$ at $p \in \iota'_t(U)$. This vector field has the following property: for any compact subset κ of S , there is an $\epsilon_\kappa > 0$ such that for all $T \in [0, 1]$ the flow $\phi_t^{V_t}(\iota'_T(\kappa))$ of V_t is well defined and satisfies $\phi_{t-T}^{V_t}(\iota'_T(\kappa)) \subset \iota'_t(\kappa)$ for all $|t - T| < \epsilon_\kappa$. We will say that any vector field satisfying this property *satisfies property Q*.

Let θ_{V_t} be the ω_M -dual of V_t . This is some closed 1-form. Because U is homotopic to S , there exists another closed 1-form ν on S such that $\nu + (\iota'_t)^*\theta_{V_t}$ is exact. Let $\tilde{\nu} = \pi^*\nu$ and let X_ν^t be a smooth family of vector fields defined only on $\iota'_t(U)$ whose ω_M -dual is $\tilde{\nu}$. We have that X_ν^t is tangent to $\iota'_t(S)$ because the fibers of π are symplectically orthogonal to S . Because X_ν^t is tangent to $\iota'_t(S)$, we have that $X_\nu^t + V_t$ satisfies property Q. Because the ω_M -dual of this vector field is exact, we have a smooth family of Hamiltonians $H_t : \iota'_t(U) \rightarrow \mathbb{R}$ whose associated Hamiltonian vector field is $X_\nu^t + V_t$. By using a cutoff function, we can assume that H_t smooth family of Hamiltonians on the whole of M .

The problem with this family of Hamiltonians is that the associated Hamiltonian vector field may not be integrable. In order to make it integrable (while still ensuring that it satisfies property Q), we do the following: Let f be some exhausting function on M . Then $f|_{\iota_t(S)}$ is also exhausting because ι_t is a proper embedding. We can perturb f by a C^0 small amount near ι_t to create a new function f_t which is equal to π^*f near $\iota_t(S) \subset \iota'_t(U)$ (here by abuse of notation we view π as the pushforward of the map $\pi : U \rightarrow S$). We can also assume that f_t is a smooth family of exhausting functions. The

Hamiltonian flow of f_t preserves $\iota_t(S)$ again because the fibers of π are orthogonal to S . Also because f_t is a small perturbation of f , we can assume that it is an exhausting function.

Choose some rapidly increasing positive function $g : \mathbb{R} \rightarrow \mathbb{R}$ such that $g \circ f_t + H_t$ is exhausting for all $t \in [0, 1]$. Because the Hamiltonian flow of $g \circ f_t$ preserves $\iota_t(S)$, we have that the flow of $g \circ f_t + H_t$ satisfies property Q. Also this Hamiltonian vector field is integrable for the following reason: Let K_t , $t \in [0, 1]$ be a smooth family of exhausting Hamiltonians. We will show that its Hamiltonian flow is integrable. Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a positive smooth function such that

$$g(x) > \sup_{t \in [0, 1], y \in K_t^{-1}(-\infty, x]} \frac{dK_t}{dt}(y).$$

We choose G to be any function with $G' = \frac{1}{g}$. Let $p(t)$ be a path in M satisfying $\frac{dp(t)}{dt} = X_{K_t}$. We have

$$\frac{d}{dt}(K_t(p(t))) = \frac{dK_t}{dt}(p(t)) + dK_t(X_{K_t}) = \frac{dK_t}{dt}(p(t)) < g(K_t(p(t))).$$

Hence

$$\frac{d}{dt}(G(K_t(p(t)))) < 1.$$

This implies that the function $t \rightarrow G(K_t(p(t)))$ is less than $t + C$ for some constant C . This means that if $p(0)$ starts in $K_0^{-1}(a)$ then $p(t)$ must be contained in $K_t^{-1}(-\infty, G^{-1}(t + C)]$. This implies that X_{K_t} is integrable.

This means that $g \circ f_t + H_t$ has an integrable Hamiltonian flow that satisfies property Q. In particular the Hamiltonian flow sends $\iota_0(S)$ to $\iota_t(S)$ at time t . $\square$

Proof. of Lemma 5.15. For induction purposes, we will assume that this Lemma is true in all dimensions less than $2n$. For $j = 1, \dots, n$, we will write A_j^t as the union $\cup_{|I|=n-j}(\cap_{i \in I} S_i^t)$. This is the union of all dimension $2j$ strata in $\cup_i S_i^t$. Suppose for $j' < j$, there is a smooth family of symplectomorphisms $\Phi_{t,j'}$ from $M \setminus A_{j'}^0$ to $M \setminus A_{j'}^t$ sending $A_{j'+1}^0 \setminus A_{j'}^0$ to $A_{j'+1}^t \setminus A_{j'}^t$. We now wish to show the same thing for j . We have (by our previous assumption) in particular a smooth family of symplectomorphisms induced by $\Phi_{t,j-1}$ sending the complement of A_{j-1}^0 to the complement of A_{j-1}^t in M . Note that $A_j^t \setminus A_{j-1}^t$ is a disjoint union of manifolds inside $M \setminus A_{j-1}^t$. All these manifolds are compactified by some smooth normal crossing divisor smoothly depending on t . Because we have assumed that the Lemma is true in all dimensions less than $2n$, there is also a smooth family of symplectomorphisms Ψ_t from $A_j^0 \setminus A_{j-1}^0$ to $A_j^t \setminus A_{j-1}^t$ starting at the identity symplectomorphism. We have a smooth family of embeddings $\iota_t := \Phi_{j-1,t}^{-1} \circ \Psi_t$ of $A_j^0 \setminus A_{j-1}^0$ into $M \setminus A_{j-1}^0$. By Lemma 5.16, there is a smooth family of Hamiltonian symplectomorphisms $F_t : M \setminus A_{j-1}^0 \rightarrow M \setminus A_{j-1}^0$ where F_0 is the identity and F_t sends the image of ι_0 to the image of ι_t . Hence

the smooth family of symplectomorphisms $\Phi_{j-1,t} \circ F_t$ sends the image of ι_0 inside $M \setminus A_{j-1}^0$ to $A_j^t \setminus A_{j-1}^t$ inside $M \setminus A_{j-1}^t$.

Hence by induction, we have found a smooth family of symplectomorphisms from $M \setminus A_{n-1}^0$ to $M \setminus A_{n-1}^t$ starting at the identity sending $A_n^0 \setminus A_{n-1}^0$ to $A_n^t \setminus A_{n-1}^t$. Because $A_{n-1}^t = S^t$ and $A_n^t \setminus A_{n-1}^t$ is the disjoint union of $S_i^t \setminus S^t$ for all i , the previous statement is the statement of the Lemma. Hence we have proven the Lemma. $\square$

Let $S_1, \dots, S_k$ be positively intersecting inside M . We can assume that $S_i \setminus \cup_{j \neq i} S_j$ is connected because we can replace a disconnected such manifold with a union of connected manifolds. Let θ be a 1 form on the complement $M \setminus (\cup_i S_i)$ such that $d\theta = \omega_M$. Let W be a small symplectic submanifold of M of dimension 2 symplectomorphic to some small disk $\mathbb{D}_\delta \subset \mathbb{C}$ of radius $\delta > 0$. Suppose that W is disjoint from S_j , $j \neq i$ and W intersects S_i orthogonally at $0 \in \mathbb{D}_\delta$ (i.e. it is tangent to the normal bundle of S_i). Let (r, ϑ) be polar coordinates on $\mathbb{D}_\delta$. Then θ pulls back to a 1-form on $W \setminus \{0\}$. We have that $d\theta = r dr \wedge d\vartheta$ on the punctured disk $\mathbb{D}_\delta \setminus \{0\}$. Hence θ is cohomologous to $\frac{r^2}{2} d\vartheta + \kappa_i d\vartheta$ for some constant κ_i . Suppose that I had some other disk W' intersecting S_i orthogonally and disjoint from the other S_j 's. Because $S_i \setminus \cup_{j \neq i} S_j$ is connected, there is a smooth family W_t joining W' and W . Hence the constant κ_i associated to W' is the same as the constant κ_i associated to W . Hence κ_i is an invariant of θ and S_i . We call κ_i the *wrapping of θ around S_i* .

Lemma 5.17. *Suppose that $S_1, \dots, S_k$ are orthogonal positively intersecting submanifolds of a compact symplectic manifold (M, ω_M) and suppose that there is a 1-form θ such that*

- (1) $d\theta = \omega_M$.
- (2) *The wrapping of θ around S_i for each i is negative.*

Then there exists a function f defined on $M \setminus \cup_i S_i$ such that $(M \setminus \cup_i S_i, \theta + df)$ has the structure of a finite type convex symplectic manifold.

The form $\theta + df$ restricted to a fiber $\mathbb{D}_\epsilon$ of the maps π_i minus the origin as described in Lemma 5.14 is $(\frac{r_i^2}{2} + \kappa_i) d\vartheta_i$ where (r_i, ϑ_i) are polar coordinates for $\mathbb{D}_\epsilon$ and $\kappa_i < 0$ is the wrapping number of θ around S_i .

Proof. of Lemma 5.17. We proceed by induction on the ordering $\prec$ mentioned in Lemma 5.14. Let $\pi_I : US_I \rightarrow S_I$ be the maps as described in the proof of Lemma 5.14. Suppose inductively we have for all $I' \prec I$ we have a function f' such that $\theta + df'$ is equal to $(\sum_{i \in I'} (\frac{r_i^2}{2} + \kappa_i) d\vartheta_i)$ on the fiber $\prod_{i \in I'} \mathbb{D}_\epsilon$. We now wish to modify f' on US_I away from a very small neighbourhood of $\cup_{j \notin I} S_j$ so that it has the properties we want.

We do this as follows: If we look near $\cup_{j \notin I} S_j$, $\theta + df'$ has the form we want because $I \cup \{j\} \prec I$ and the map π_I inside $US_{I \cup \{j\}}$ preserves the fibers

of $\pi_{I \cup \{j\}}$ and corresponds to the projection

$$\prod_{i \in I \cup \{j\}} \mathbb{D}_\epsilon \twoheadrightarrow \prod_{i \in I} \mathbb{D}_\epsilon.$$

eliminating the $\mathbb{D}_\epsilon$ corresponding to j . Because $\theta + df'$ restricted to a fiber of π_I is cohomologous to $\sum_{i \in I} (\frac{r_i^2}{2} + \kappa_i) d\vartheta_i$, there is a smooth function f'' from US_I to $\mathbb{R}$ such that $\theta + df' + df''$ restricted to each fiber is of the form $\sum_{i \in I} (\frac{r_i^2}{2} + \kappa_i) d\vartheta_i$. Such a function exists because $\theta + df' - \sum_{i \in I} (\frac{r_i^2}{2} + \kappa_i) d\vartheta_i$ restricted to each fiber of π_I is exact, so for each fiber $\pi_I^{-1}(q)$ we have a function f_q which is unique up to adding a constant such that this 1-form is equal to df_q . We can assume that f_q smoothly depends on q by adding some function of the form $\kappa \circ \pi_I$ (this basically follows from a parameterized version of the Poincaré Lemma). Hence we can view f_q as a function f'' which has the properties we want.

Near $\cup_{j \notin I} S_j$ we have that $\theta + df'$ already has the form we want, hence f'' must be of the form $\pi_I^* h$ in this region. By extending h to the whole of S_I , we can subtract $\pi_I^* h$ from f'' so that $f'' = 0$ near $\cup_{j \notin I} S_j$. This means that $\theta + df' + df''$ restricted to each fiber of π_I is equal to $\sum_{i \in I} (\frac{r_i^2}{2} + \kappa_i) d\vartheta_i$, and also because $f'' = 0$ near $\cup_{j \notin I} S_j$, it also has this form for all other maps $\pi_{I'}$ where $I' \prec I$. Hence by induction we have shown that there is some function f such that $\theta + df = \sum_{i \in I} (\frac{r_i^2}{2} + \kappa_i) d\vartheta_i$ inside the fibers of π_I for all I . This is the second part of the Lemma.

We now wish to show that we have the structure of a finite type convex symplectic manifold. Let $\nu : (0, \epsilon) \rightarrow \mathbb{R}$ be a function which is equal to 0 near ϵ and tends to infinity as we reach 0 and which has non-positive derivative. We assume it has negative derivative near 0. Then we view $\nu(r_i)$ as a function on the complement of S_i as we can extend it by zero away from US_i . We have that $\sum_i \nu(r_i)$ is an exhausting function. Let $X_{\nu(r_i)}$ be the Hamiltonian vector field of $\nu(r_i)$. Because the ω -orthogonal plane bundle to the fibers $\mathbb{D}_\epsilon$ of US_i are contained in the level sets of r_i , we have that $X_{\nu(r_i)}$ is tangent to the fibers of π_i inside US_i and is zero elsewhere. Also $X_{\nu(r_i)}$ restricted to some fiber $\mathbb{D}_\epsilon$ is equal to $X_{\nu(r_i)|_{\mathbb{D}_\epsilon}}$ which is equal to $-\frac{\nu'(r_i)}{r_i} \frac{\partial}{\partial \vartheta_i}$. Hence $X_{\nu(r_i)}(\theta + df) < 0$ for r small enough because κ_i is negative. If $X_{\theta+df}$ is the ω_M -dual of $\theta + df$ then $X_{\theta+df}(d(\sum_i \nu(r_i))) = -(\sum_i X_{\nu(r_i)})(\theta + df) \geq 0$. This is greater than zero near $\cup_i S_i$ because for each point p near $\cup_i S_i$, we have that $\nu'(r_i) > 0$ for some i . Hence $(M \setminus \cup_i S_i, \theta + df)$ has the structure of a finite type convex symplectic manifold. This proves the Lemma. $\square$

Let P be a smooth projective variety, and D some effective divisor so that

- (1) D is a smooth normal crossing divisor. i.e. it is a union of smooth complex hypersurfaces which are transversally intersecting.
- (2) $P \setminus D$ is isomorphic to A where A is a smooth affine variety.

Let L be a line bundle on P and $\|\cdot\|$ a metric on L so that if F is its curvature form then $-iF$ restricted to A is a Kähler form but this may not

be true along D . Let s be a section of L so that $s^{-1}(0) = D$. We will call $d^c \log \|s\|$ a *compactification 1-form* associated to A . Recall that we have another 1-form on A coming from an embedding $\iota : A \hookrightarrow \mathbb{C}^N$. This is described before Lemma 2.1. This is given by $\iota^* \sum_i \frac{r_i^2}{2} d\vartheta_i$ where (r_i, θ_i) are polar coordinates on the i th factor of $\mathbb{C}^N$. The following lemma is almost exactly the same as [Sei08, Lemma 4.4].

Lemma 5.18. *We have that $(A, d^c \log \|s\|)$ makes A into a finite type convex symplectic manifold convex deformation equivalent to (A, θ_A) .*

Proof. of Lemma 5.18. This lemma will be done in three steps. In step 1 we show that $(A, d^c \log \|s\|)$ is a finite type convex symplectic manifold. In step 2 we will show that if $d^c \log \|s'\|'$ is another compactification 1-form then $(A, d^c \log \|s\|)$ is convex deformation equivalent to $(A, d^c \log \|s'\|')$. Finally in step 3 we will show that $(A, d^c \log \|s\|)$ is convex deformation equivalent to (A, θ_A) .

Step 1: Let $p \in D$ and choose local holomorphic coordinates $z_1, \dots, z_n$ and a trivialization of L around p so that $s = z_1^{w_1} \cdots z_n^{w_n}$ ($w_i \geq 0$). The metric $\|\cdot\|$ on L is equal to $e^\psi |\cdot|$ for some function ψ with respect to this trivialization where $|\cdot|$ is the standard metric on $\mathbb{C}$. So

$$d \log \|s\| = -\psi - \left(\sum_i w_i d \log |z_i| \right).$$

If we take the vector field $Y := -z_1 \partial_{z_1} \cdots - z_n \partial_{z_n}$, then $d \log (|z_j|)(Y) = -1$ and $Y \cdot \psi$ tends to zero. Hence $d \log \|s\|$ is non-zero near infinity. If $X_{d^c \log \|s\|}$ is the $dd^c \log \|s\|$ -dual of $d^c \log \|s\|$ then $-d \log (\|s\|)(X_{d^c \log \|s\|}) = \|d \log \|s\|\|^2 > 0$ near infinity. Hence $(A, d^c \log \|s\|)$ is a finite type convex symplectic manifold.

Step 2: Suppose now that $P', D', L', s', \|\cdot\|'$ are different choices of compactification, effective divisor, line bundle, section and metric satisfying similar properties to P, D, L, s and $\|\cdot\|$. Here $P' \setminus D'$ is isomorphic to A . By applying the Hironaka resolution of singularities theorem again [Hir64] we have a third compactification P'' and morphisms $\pi : P'' \rightarrow P$, $\pi' : P'' \rightarrow P'$. We pull back L and L' to P'' as well as the sections s, s' giving us new line bundles $\tilde{L}, \tilde{L}'$ and sections $\tilde{s}$ and $\tilde{s}'$. By abuse of notation we write $\|\cdot\|$ and $\|\cdot\|'$ for the metrics on $\tilde{L}$ and $\tilde{L}'$ which are the pullbacks of $\|\cdot\|$ and $\|\cdot\|'$ respectively. Again we look at some local coordinate chart $z_1, \dots, z_n$ and vector field Y around some point $p \in D''$ where $D'' = P'' \setminus A$. Using the same arguments as before we have

$$(t \log \|\tilde{s}\| + (1-t) \log \|\tilde{s}'\|')(Y) > 0$$

near infinity for all $t \in [0, 1]$. Hence $(A, d^c \log \|s\|)$ and $(A, d^c \log \|s'\|')$ are convex deformation equivalent.

Step 3: In this step we will need to refer to the proof of Lemma 2.1 so the reader must be familiar with this lemma first. Let $R := \sum_i \frac{r_i^2}{4}$. We have that $\theta_A = -d^c R$. Basically from the proof of Lemma 2.1 we have

a compactification 1-form equal to $-d^c f(R)$ where f is a non-decreasing function. We have that $(A, -d^c(tR + (1-t)f(R)))$ is a convex deformation equivalence from (A, θ_A) to $(A, -d^c f(R))$. The reason why this is a convex deformation equivalence is that the level sets of R and $f(R)$ coincide (up to reparameterization). By Step 2 we have (A, θ) is convex deformation equivalent to $(A, -d^c f(R))$. Hence (A, θ) is convex deformation equivalent to (A, θ_A) . $\square$

Lemma 5.19. *Let M be a smooth projective variety and let $\cup_i S_i$ be a smooth normal crossing divisor such that $\sum_i a_i S_i$ is ample for some $a_1, \dots, a_k \in \mathbb{N} \setminus \{0\}$. The complement $A := M \setminus \cup_i S_i$ is an affine variety and it has a natural 1-form θ with $d\theta = \omega_M$. We have that (A, θ) is convex deformation equivalent to A with the natural 1-form θ_A coming from some embedding of A into $\mathbb{C}^N$ (see Section 2.1). Then $S_1, \dots, S_k$ are positively intersecting and the wrapping number of θ around S_i is negative.*

Proof. of Lemma 5.19. The natural 1-form θ is a compactification 1-form as described earlier. We have an ample line bundle L coming from $\sum_i a_i S_i$. There is a metric $\|\cdot\|$ on L whose associated curvature form F has the property that iF is a positive $(1,1)$ -form. This is our symplectic form ω_M on M . Choose a holomorphic section s of L whose zero set is exactly $\sum_i a_i S_i$ (i.e. the zero set has multiplicity a_i at S_i). Then our 1-form θ is $d^c \log \|s\|$ where $d^c = d \circ J$ where J is our complex structure on M .

First of all, the divisors S_i are positively intersecting submanifolds because J is compatible with the symplectic form ω_M and all the S_i 's are holomorphic submanifolds. So the hard part is proving that θ wraps negatively around S_i . Let p be a point in S_i disjoint from $\cup_{j \neq i} S_j$ and choose a small holomorphic disk $\mathbb{D}_\delta$ disjoint from $\cup_{j \neq i} S_j$ and intersecting S_i at p only. We can assume that this disk is intersecting S_i orthogonally because the ω_M -orthogonal bundle to S_i is holomorphic. The line bundle L restricted to this disk $\mathbb{D}_\delta \subset \mathbb{C}$ is equal to $a_i \{0\}$ where $\{0\}$ means the divisor at 0 and the section s_i has a zero of multiplicity i at 0. This means we can choose a trivialization of L so that s_i is equal to z^{a_i} where z is the complex coordinate on $\mathbb{D}_\delta$.

We have that $dd^c \log \|z^{a_i}\| = dd^c \log \|1\|$ so the wrapping number of θ around S_i corresponds to the cohomology class of

$$\begin{aligned} d^c \log \|z^{a_i}\| - d^c \log \|1\| &= a_i d^c \log |z| + d^c \log \|1\| - d^c \log \|1\| \\ &= a_i d^c \log |z| = -a_i d\vartheta. \end{aligned}$$

Here $|\cdot|$ is the standard euclidean metric on $\mathbb{D}_\delta \subset \mathbb{C}$ and ϑ is the angular coordinate on $\mathbb{D}_\delta$. This is a negative multiple of $d\vartheta_i$ and hence we have that θ wraps around S_i negatively. By Lemma 5.18 we get that (A, θ_A) is convex deformation equivalent to (A, θ) . $\square$

Putting all these Lemmas together gives us the following theorem:

Theorem 5.20. *Let A be a smooth affine variety. Then A is convex deformation equivalent to a finite type convex symplectic manifold (W, θ_W) with the following properties:*

- (1) W is symplectomorphic to $M \setminus \cup_i S_i$ where S_i are co-dimension 2 symplectic submanifolds transversely intersecting.
- (2) There are neighborhoods US_i of S_i and fibrations $\pi_i : US_i \twoheadrightarrow S_i$ satisfying the properties stated in Lemma 5.14.
- (3) θ_W restricted to the fiber $\mathbb{D}_\epsilon$ of π_i is equal to $(r_i^2 + \kappa_i)d\vartheta_i$ for some $\kappa_i < 0$.

Proof. of Theorem 5.20. We first compactify A to a smooth projective variety and then by using [Hir64], we blow up this projective variety so that $A = M \setminus \cup_i S'_i$ for some transversely intersecting complex hypersurfaces S'_i inside a new smooth projective variety M . These are positively intersecting symplectic submanifolds of M and the natural 1-form θ ($d\theta = \omega_M$) on A wraps negatively around S_i for each i by Lemma 5.19.

By Lemma 5.3, there is a smooth family of positively intersecting symplectic submanifolds S_i^t such that $S_i^0 = S'_i$ and S_i^1 are orthogonal. Let $S^t := \cup_{i,j} (S_i^t \cap S_j^t)$. By Lemma 5.15, there is a smooth family of symplectomorphisms Φ_t from $M \setminus S^0$ to $M \setminus S^t$ starting at the identity and sending $S_i^0 \setminus S^0$ to $S_i^t \setminus S^t$.

This means that the symplectomorphisms Φ_t also induce symplectomorphisms from $M \setminus \cup_i S_i^0$ to $M \setminus \cup_i S_i^t$. Also $\theta_t := (\Phi_t|_{M \setminus \cup_i S_i^0})_* \theta$ wraps around S_i^t negatively because the symplectomorphism $\Phi_t|_{M \setminus \cup_i S_i^0}$ extends smoothly over S_i to Φ_t . This means by Lemma 5.17, there is an exact 1-form df such that $\theta_1 + df$ makes $M \setminus \cup_i S_i^1$ into a finite type convex symplectic manifold with the property that $\theta_1 + df$ restricted to each fiber $\mathbb{D}_\epsilon$ of π_i is equal to $(r_i^2 + \kappa_i)d\vartheta_i$ where $\kappa_i < 0$. So if we set $W := M \setminus \cup_i S_i^1$ and $\theta_W := \theta_1 + df$ then we have proven the Lemma. $\square$

6. THE GROWTH RATE OF AFFINE VARIETIES

Definition 6.1. Let $(M_1, \theta_1), (M_2, \theta_2)$ be two exact symplectic manifolds. We say that M_1, M_2 are exact symplectomorphic at infinity if there are open subsets $U_i \subset M_i$ and a symplectomorphism

$$\Psi : M_1 \setminus U_1 \rightarrow M_2 \setminus U_2$$

such that

- (1) The closure of U_i is compact inside M_i .
- (2) Ψ sends compact sets in $M_1 \setminus U_1$ to compact sets in $M_2 \setminus U_2$.
- (3) Ψ is an exact symplectomorphism (i.e. $\Psi^* \theta_2 = \theta_1 + df$).

Definition 6.2. Let $(W_1, \theta_{W_1}), (W_2, \theta_{W_2})$ be two convex symplectic manifolds. We say that they are convex deformation equivalent at infinity if there is a sequence of convex symplectic manifolds (Q_i, θ_{Q_i}) for $i = 1, \dots, k$ such that:

- (1) Q_1 is convex deformation equivalent to W_1 .
- (2) Q_k is convex deformation equivalent to W_2 .

- (3) Q_i is convex deformation equivalent to Q_{i+1} or they are exact symplectomorphic at infinity.

Note that being *convex deformation equivalent at infinity* is an equivalence relation.

Theorem 6.3. *Suppose that a finite type convex symplectic manifold B and a smooth affine variety A are convex deformation equivalent at infinity. Then filtered directed system $(SH_*^\#(B, \theta_B, \lambda))$ is isomorphic to a filtered directed system (V_x) where the dimension $|V_x|$ satisfies $|V_x| \leq P(x)$ for some polynomial P . The degree of this polynomial is less than or equal to m_A where m_A is defined in the introduction.*

This has the following direct corollary:

Theorem 6.4. *We have $\Gamma(B) \leq m_A$.*

Theorem 1.1 follows directly from Theorem 6.4 where we set $B = A$. Theorem 1.2 also follows from Corollary 6.3. Here is a statement of this theorem: *Suppose that the boundary of a Liouville domain M is algebraically Stein fillable, then $\Gamma(M) \leq m_A$.*

Proof. of Theorem 1.2. The boundary of M is contactomorphic to the boundary of $\bar{A}$. This means that we can deform M through a family of Liouville domains to a new Liouville domain M' such that the contact form on $\partial M'$ coincides exactly with the contact form on $\partial \bar{A}$. This means that the completions $\widehat{M}'$ and $A := \widehat{\bar{A}}$ are exact symplectomorphic at infinity because their cylindrical ends are identical. Also $\widehat{M}'$ is convex deformation equivalent to $\widehat{M}$ which means that $\widehat{M}$ and $\widehat{A}$ are convex deformation equivalent at infinity. By Theorem 6.4, we have that $\Gamma(\widehat{M}) < \infty$ which proves Theorem 1.2. $\square$

Before we prove Theorem 6.3, We need some lemmas: The first Lemma gives us a slightly more general definition of growth rate which will be useful for our purposes. Let (H, J) be a growth rate admissible Hamiltonian on some finite type convex symplectic manifold (W, θ_W) and let $\kappa > 0$ be any constant. Let $\lambda_1 \leq \lambda_2$. Then there is a natural morphism from $SH_*^{[0, \kappa \lambda_1]}(\lambda_1 H, J)$ to $SH_*^{[0, \kappa \lambda_2]}(\lambda_2 H, J)$ given by first composing the morphism

$$SH_*^{[0, \kappa \lambda_1]}(\lambda_1 H, J) \rightarrow SH_*^{[0, \kappa \lambda_1]}(\lambda_2 H, J)$$

induced by the continuation map followed by the morphism

$$SH_*^{[0, \kappa \lambda_1]}(\lambda_2 H, J) \rightarrow SH_*^{[0, \kappa \lambda_2]}(\lambda_2 H, J)$$

given by the natural inclusion map of chain complexes. We could have done this the other way around by starting with an inclusion of chain complexes:

$$SH_*^{[0, \kappa \lambda_1]}(\lambda_1 H, J) \rightarrow SH_*^{[0, \kappa \lambda_2]}(\lambda_1 H, J)$$

and then composing it with a continuation map. But this gives us exactly the same map. Because we can swap inclusion maps and continuation maps,

we have that these maps satisfy functoriality properties which means we get a filtered directed system:

$$(SH_*^{[0, \kappa\lambda]}(\lambda H, J))$$

Lemma 6.5. *The filtered directed system described above is isomorphic to $(SH_*^\#(W, \theta_W))$.*

Proof. of Lemma 6.5. We have a natural map ϕ from $SH_*^{[0, \kappa\lambda]}(\lambda H, J)$ to $SH_*^\#(\lambda H, J)$ given by inclusion of chain complexes. This commutes with the filtered directed system maps so is a morphism of filtered directed systems. There is a constant $K > 1$ and a 1-form θ such that $\theta - \theta_W$ is exact, $\kappa K > 1$ and $-\theta(X_H) - H < \kappa K$ by the action bound property. We have a map ϕ' from $SH_*^\#(\lambda H, J)$ to $SH_*^{[0, \kappa K\lambda]}(K\lambda H, J)$ constructed as follows: First of all we have an isomorphism from $SH_*^\#(\lambda H, J)$ to $SH_*^{[0, \kappa K\lambda]}(\lambda H, J)$ because all the 1-periodic orbits have action less than $\kappa K\lambda$. We then compose this isomorphism with the continuation map:

$$SH_*^{[0, \kappa K\lambda]}(\lambda H, J) \rightarrow SH_*^{[0, \kappa K\lambda]}(K\lambda H, J).$$

Again this commutes with the filtered directed system maps because continuation maps and action inclusion maps commute.

The map $\phi \circ \phi'$ is the composition:

$$\begin{aligned} SH_*^\#(\lambda H, J) &\rightarrow SH_*^{[0, \kappa K\lambda]}(\lambda H, J) \rightarrow \\ SH_*^{[0, \kappa K\lambda]}(K\lambda H, J) &\rightarrow SH_*^\#(K\lambda H, J). \end{aligned}$$

This is the natural continuation map because action inclusion and continuation maps commute so the above composition is equal to the following composition:

$$\begin{aligned} SH_*^\#(\lambda H, J) &\rightarrow SH_*^{[0, \kappa K\lambda]}(\lambda H, J) \rightarrow \\ SH_*^\#(\lambda H, J) &\rightarrow SH_*^\#(K\lambda H, J). \end{aligned}$$

Also because all the 1-periodic orbits of λH have action less than $\kappa K\lambda$, we have that the composition of the first two maps is the identity map. Hence $\phi \circ \phi'$ is a continuation map.

We also have that $\phi' \circ \phi$ is a continuation map because it is equal to the natural composition:

$$\begin{aligned} SH_*^{[0, \kappa\lambda]}(\lambda H, J) &\rightarrow SH_*^\#(\lambda H, J) \rightarrow \\ SH_*^{[0, K\kappa\lambda]}(\lambda H, J) &\rightarrow SH_*^{[0, K\kappa\lambda]}(K\lambda H, J). \end{aligned}$$

Because all 1-periodic orbits of λH have action less than $\kappa K\lambda$ the composition of the first two maps is identical to the natural inclusion map. Hence $\phi' \circ \phi$ is a directed system map and so ϕ, ϕ' give us our isomorphism of filtered directed systems. $\square$

Lemma 6.6. *Suppose we have a Hamiltonian $H : W \rightarrow \mathbb{R}$, a function $P : \mathbb{R} \rightarrow \mathbb{R}$ and a small open neighbourhood $\mathcal{N}$ of $H^{-1}(0)$ such that:*

- (1) H satisfies the Liouville vector field property.
- (2) For every λ outside some discrete subset, there is a C^2 small perturbation H_λ of λH such that all the 1-periodic orbits of H_λ inside $\mathcal{N}$ are non-degenerate and the number of such orbits is bounded above by $P(\lambda)$.

Then there is a filtered directed system (V_λ) isomorphic to $(SH_*^\#(W, \theta_W))$ such that the rank of V_λ is bounded above by $P(\lambda)$.

Proof. of Lemma 6.6. If H was growth rate admissible and $\mathcal{N} = W$ then this lemma would be fairly straightforward as the rank of SH_* is bounded above by the number of non-degenerate orbits. The problem is that it may not be growth rate admissible. Instead we will construct a growth rate admissible Hamiltonian H^p such that all the 1-periodic orbits of sufficiently low action are the same as the ones of H and then invoke Lemma 6.5.

By Lemma 4.2, there is a growth rate admissible pair (H^p, J^p) such that

- (1) $H^p = H$ on a small neighbourhood of $H^{-1}(0)$ and $(H^p)^{-1}(0) = H^{-1}(0)$.
- (2) $-\theta_H(X_{H^p}) - H^p \geq 0$ everywhere.
- (3) $-\theta_H(X_{H^p}) - H^p > 0$ when $H^p > 0$.
- (4) $-\theta_H(X_{H^p}) - H^p$ is greater than some constant $\delta_H > 0$ outside some compact set.

Here θ_H is the 1-form that makes H satisfy the Liouville vector field property.

Let A be the function $-\theta_H(X_{H^p}) - H^p$. The level sets of H^p near $(H^p)^{-1}(0)$ are compact because H^p satisfies the bounded below property. Let $\nu_H > 0$ be a constant so that for all $x \in (0, \nu_H]$ we have that $(H^p)^{-1}(x)$ is compact and regular and contained inside $\mathcal{N} \cap \{H^p = H\}$. Also because $A(x) > \delta_H$ outside some compact set, $A \geq 0$ and $A(y) > 0$ if and only if $H^p(y) > 0$, there is a constant $\epsilon_A > 0$ with the property that $A^{-1}([0, \epsilon_A])$ is contained in the region $(H^p)^{-1}([0, \nu_H])$. For $\lambda \geq 1$, any 1-periodic orbit of λH^p of action $\leq \lambda \epsilon_A$ must be contained in the region $(H^p)^{-1}([0, \nu_H])$. This orbit is also contained inside $\mathcal{N}$ and the region where H^p is equal to H .

Choose a sequence of perturbations K_λ^i that C^2 converge to λH where the number of 1-periodic orbits of K_λ^i contained inside $\mathcal{N}$ is bounded above by $P(\lambda)$ and so that all of these orbits are non-degenerate. We make the perturbation small enough so that orbits of action less than or equal to $\lambda \epsilon_A$ are contained in some fixed open subset U of $\{H^p = H\} \cap \mathcal{N}$ such that U contains $(H^p)^{-1}([0, \epsilon])$. Let $\rho : W \rightarrow \mathbb{R}$ be a bump function which is 0 on a small neighbourhood of the closure of U and 1 in the region where $H^p \neq H$. Let $H_\lambda^i := (1 - \rho)K_\lambda^i + \rho\lambda H^p$. These Hamiltonians C^1 converge to λH^p . Hence for large enough i , we have that the action of all the 1-periodic orbits of H_λ^i that are not entirely contained in U is greater than $\frac{\lambda \epsilon_A}{2}$. For large enough i we also have a sequence of constants δ_i tending to 0 such that $(H_\lambda^i)^{-1}([0, \lambda \epsilon - \delta_i])$ contains all the orbits of action $\leq \frac{\lambda \epsilon_A}{2}$ and so that there are no orbits on the boundary of $U_i := (H_\lambda^i)^{-1}([0, \lambda \epsilon_A - \delta_i])$. We can also assume that $U_i \subset U$.

By Lemma 2.2, we perturb H_λ^i again to $H_\lambda'^i$ by a C^2 small amount outside the closed set U_i so that $H_\lambda'^i = K_\lambda^i$ inside U_i , all of its 1-periodic orbits outside U_i are non-degenerate of action greater than $\frac{\lambda\epsilon_A}{2}$ and such that it is equal to H_λ^i near infinity. For large enough i , $(H_\lambda'^i, J^p)$ is growth rate admissible where H_λ^i has only non-degenerate orbits and such that all of its orbits of action less than or equal to $\frac{\lambda\epsilon_A}{2}$ are in the region where this Hamiltonian is equal to K_λ^i . Hence the number of 1-periodic orbits of $H_\lambda'^i$ of action less than or equal to $\frac{\lambda\epsilon_A}{2}$ is bounded above by $P(\lambda)$. This implies that the rank of $V_\lambda := SH_*^{[0, \frac{\lambda\epsilon_A}{2}]}(H_\lambda'^i, J^p)$ is bounded above by $P(\lambda)$. By Lemma 6.5, the filtered directed system (V_λ) is isomorphic to $SH_*^\#(W, \theta_W)$. This proves the Lemma. $\square$

Suppose we have a Hamiltonian H and a function P such that there is a small neighbourhood N of $H^{-1}(0)$ where H, P, N satisfy the properties of the previous Lemma, then we say that H is P bounded.

Lemma 6.7. *Suppose that (W, θ_W) and $(W', \theta_{W'})$ are finite type convex symplectic manifolds that are convex deformation equivalent at infinity. Suppose also that we have a Hamiltonian H on W and a function $P : \mathbb{R} \rightarrow \mathbb{R}$ such that:*

- (1) H satisfies the Liouville vector field property.
- (2) H is P bounded.

Then W' also admits a Hamiltonian H' satisfying the Liouville vector field property and that is P bounded.

Proof. of Lemma 6.7. Let X_{θ_W} be the $d\theta_W$ -dual of θ_W . Let f_W be an exhausting function such that $df_W(X_{\theta_W}) > 0$ when $f_W \geq C_W$ for some C_W . By Lemma 8.1, there is a family of 1-forms θ_W^s such that:

- (1) $d\theta_W^s$ is symplectic.
- (2) $\theta_W = \theta_W^0$.
- (3) $\theta_W^s = \theta_W$ in the region $f^{-1}(-\infty, C_W]$.
- (4) θ_W^1 is complete.
- (5) The $d\theta_W^s$ -dual $X_{\theta_W^s}$ of θ_W^s satisfies $X_{\theta_W^s} = g_s X_{\theta_W^0}$ for some positive function g_s which smoothly depends on s .

We have a similar family of 1-forms $\theta_{W'}^s$ on W' and a similar function $f_{W'}$ such that these 1-forms are all equal on $f_{W'}^{-1}(-\infty, C_{W'}]$. The pairs (W, θ_W^s) (resp. $(W', \theta_{W'}^s)$) are all finite type convex symplectic manifolds because their Liouville vector fields are all of the form $g_s X_{\theta_W}$ (resp. $h_s X_{\theta_{W'}}$) for some smooth family of functions g_s (resp. h_s) and the same reason ensures that this is a convex deformation. We can assume that the closure of $H^{-1}(0)$ is contained in $f_W^{-1}(-\infty, C_W)$. Hence $\theta_W = \theta_W^s$ on some compact set whose interior contains $H^{-1}(0)$. This implies that H satisfies the Liouville vector field property and is P bounded on (W, θ_W^s) for all s and in particular for (W, θ_W^1) . Let $X_{\theta_W^1}$ be the $d\theta_W^1$ -dual of θ_W^1 . If Φ_t is the flow of this vector

field then $\Phi_t^* \theta_W^1 = e^t \theta_W^1$ which implies that $H_t := \Phi_t^*(H)$ is still P bounded. In particular we can ensure that the compact set $H_t^{-1}(0)$ is arbitrarily large.

Because (W, θ_W^1) and $(W', \theta_{W'}^1)$ are complete finite type convex symplectic manifolds that are convex deformation equivalent at infinity, we have by Lemma 8.7 that they are exact symplectomorphic at infinity. Let $\phi : W \setminus K_W \rightarrow W' \setminus K_{W'}$ be this exact symplectomorphism. By the previous discussion, we can assume that the interior of $H_t^{-1}(0)$ contains K_W . We define K' on W' by pushing H_t forward via ϕ and then defining K' to be zero inside $K_{W'}$. This Hamiltonian still satisfies the Liouville vector field property and is P bounded on $(W', \theta_{W'}^1)$. Let Φ'_t be the flow of the vector field $X_{\theta_{W'}^1}$. For large enough T , $H' := (\Phi'_T)^* K'$ has the property that $(H')^{-1}(0)$ is a subset of $f_{W'}^{-1}(-\infty, C_{W'})$. Because $\theta_W^s = \theta_{W'}^s$ inside $f_{W'}^{-1}(-\infty, C_{W'})$, we have that H' is P bounded on $(W', \theta_{W'}^s)$ for all s and in particular on $(W', \theta_{W'})$. $\square$

Given some smooth affine variety A , we will construct a Hamiltonian H on A satisfying the Liouville vector field property. By Theorem 5.20, the smooth affine variety A is exact symplectomorphic to a finite type convex symplectic manifold W described as follows: We start with a compact symplectic manifold M and a set of transversely intersecting symplectic submanifolds S_i ($i = 1, \dots, k$). We write $S_{\{i_1, \dots, i_l\}}$ for the intersection $\cap_j S_{i_j}$. There are small neighborhoods US_i of S_i and projections $\pi_i : US_i \rightarrow S_i$ and a positive integer n_W (the number of such S_i 's) such that

- (1) For $1 \leq i_1 < i_2 < \dots < i_l \leq n_W$,

$$\pi_{i_l} \circ \dots \circ \pi_{i_1} : \cap_{j=1}^l US_{i_j} \rightarrow S_{\{i_1, \dots, i_l\}}$$

has fibers that are symplectomorphic to $\Pi_{j=1}^l \mathbb{D}_\epsilon$ where $\mathbb{D}_\epsilon$ is the disk of radius ϵ .

- (2) If we look at a fiber $\Pi_{j=1}^l \mathbb{D}_\epsilon$ of $\pi_{i_l} \circ \dots \circ \pi_{i_1}$, then for $1 \leq m \leq l$, π_{i_m} maps this fiber to itself. It is equal to the natural projection

$$\Pi_{j=1}^l \mathbb{D}_\epsilon \rightarrow \Pi_{j=1, j \neq m}^l \mathbb{D}_\epsilon$$

eliminating the m th disk $\mathbb{D}_\epsilon$.


- (3) The symplectic structure on US_i induces a natural connection for $\pi_{i_l} \circ \dots \circ \pi_{i_1}$ given by the ω orthogonal vector bundle to the fibers. We require the associated parallel transport maps to be elements of $U(1) \times \dots \times U(1)$ where $U(1)$ acts on the disk $\mathbb{D}_\epsilon$ by rotation.

The finite type convex symplectic manifold W we want is symplectomorphic to $M \setminus \cup_i S_i$. Let $r_i : US_i \rightarrow \mathbb{R}$ be the function such that when we restrict r_i to a fiber D_ϵ of π_i , we get that r_i is the distance from the origin. This uniquely determines r_i because π_i has a $U(1)$ structure group.

We will now construct our Hamiltonian on the Liouville manifold W . Let $\nu : [0, \epsilon) \rightarrow [0, \infty)$ be a smooth function satisfying:

- (1) Near $t = \epsilon$, $\nu(t) = 0$ and near 0, $\nu(t) = \frac{\epsilon^2}{4} - t^2$.

- (2) $\nu' < 0$ when $\nu > 0$.
- (3) There is one point x where $\nu''(x) = 0$ and $\nu(x) > 0$ (i.e. the graph of ν has exactly one point of inflection). In particular, $\nu''(x)$ is negative when $\nu(x) > 0$ is small.
- (4) If $\nu(x) > 0$ is small then we require that $x < -\kappa_i$ for each i and $x < 1$.



The function $\nu(r_i) : US_i \rightarrow \mathbb{R}$ can be extended smoothly by 0 to the whole of M . From now on we will write $\nu(r_i)$ as this function from M to $\mathbb{R}$. Let $H := \sum_i \nu(r_i)$. Theorem 5.20 also says that the Liouville form θ_W on W satisfies: θ_W restricted to a fiber $\mathbb{D}_\epsilon$ of π_i is equal to $(r_i^2 + \kappa_i)d\vartheta_i$ where r_i, ϑ_i are polar coordinates on $\mathbb{D}_\epsilon$ and $\kappa_i < 0$ is some constant. Because the level sets of $\nu(r_i)$ contain the ω_A -orthogonal vector bundle to the fibers of π_i , we have that its Hamiltonian flow is contained in the fibers and is equal to $-\frac{1}{2r_i}\nu'(r_i)\frac{\partial}{\partial \vartheta_i}$. We have that $\theta_W(X_H) = -\frac{r_i^2 + \kappa_i}{2r_i}\nu'(r_i)$. Hence if X_{θ_W} is the $d\theta_W$ -dual of θ_W then $dH(X_{\theta_W}) = -\theta_W(X_H) \geq 0$ and $dH(X_{\theta_W}) = -\theta_W(X_H) > 0$ when $H > 0$. The derivative with respect to r_i of $\theta_W(X_H)$ is $\left(-\frac{1}{2} + \frac{\kappa_i}{2r_i^2}\right)\nu'(r_i) - \frac{r_i^2 + \kappa_i}{2r_i}\nu''(r_i)$ which is negative when $\nu(r_i) > 0$ is small because $\nu''(r_i) \gg \nu'(r_i)$ and κ_i is negative. Hence $d(dH(X_{\theta_W}))(X_{\theta_W}) = \theta_W(X_{\theta_W(X_H)}) > 0$. Hence H satisfies the Liouville vector field property. We will call H a *Hamiltonian compatible with a compactification of A* .

Lemma 6.8. *The Hamiltonian H is P bounded for some polynomial of degree at least d where*

$$d = n - \min\left(\frac{1}{2}\dim_{\mathbb{R}}\{S_I | S_I \neq \emptyset\}\right).$$

Proof. of Lemma 6.8. The basic idea here is that the orbits of λH form manifolds with corners coming from the divisors. The number of such manifolds is bounded above by some constant times λ^d and the number of such manifolds up to diffeomorphism is finite. We use a Morse function on each of these manifolds with corners to perturb them into non-degenerate orbits and this gives us our bound. From now on we will assume that W is connected.

We write US_I for $\cap_{i \in I} US_i$. The Hamiltonian flow of λH in the region $US_I \setminus \cup_{J \subsetneq I} US_J$ fixes the fibers $\prod_{i \in I} \mathbb{D}_\epsilon$ and for each $i \in I$ it rotates the i 'th fiber in this product by an angle of $-\lambda \frac{1}{2r_i} \nu'(r_i)$. Hence the fixed points of the time- λ flow H form manifolds with corners. These are diffeomorphic to $\cap_{i \in I} US_i$ minus the union of the interiors of US_i for all $i \notin I$. There is also one such connected manifold of co-dimension 0 which is diffeomorphic to W minus the union of the interiors of US_i for all i . Let τ be a constant greater than the supremum of $-\frac{4\nu'(t)}{t}$. This is bounded because $\nu'(t)$ is a multiple of t near 0. There are at most $\lfloor \tau \lambda / 2\pi \rfloor^m \binom{n_w}{m}$ such manifolds of codimension m for $1 \leq m \leq d$ and 0 such manifolds for $m > d$. Also there are at most $\sum_{m=0}^d \binom{n_w}{m}$ diffeomorphism types of such manifolds.

Let $h : M \rightarrow \mathbb{R}$ be a function such that h restricted to $\cap_j S_{i_j}$ is Morse for any $i_1 < i_2 < \dots < i_l$. We also assume it has the following properties:

- (1) h is a C^∞ small perturbation of H .
- (2) On a small neighbourhood of the closure of $\nu(r_i) > 0$, we have that $h = \nu(r_i(1 - \delta_h))$ for some small $\delta_h > 0$.

We now perturb λH as follows: Consider a manifold Y of fixed points of the Hamiltonian flow of λH in the region where there is an $i_1 < i_2 < \dots < i_l$ such that $\nu'(r_k) \neq 0$ if and only if $k = i_j$ for some j . This manifold is contained in the region $\cap_j US_{i_j}$. Let N_Y be a small neighbourhood of Y such that the only fixed points of λH inside N_Y are ones in Y . The manifold is of co-dimension l and is a co-dimension 0 submanifold with corners inside the closed manifold $\bar{Y} := \cap_{j=1}^l \{r_{i_j} = s_j\}$ for some $s_1, \dots, s_l \in \mathbb{R}$. In fact $Y = \bar{Y} \cap \cap_{k \neq i_j \forall j} \{\nu(r_k) = 0\}$. Let $\rho : N_Y \rightarrow \mathbb{R}$ be a bump function which is 0 outside some compact set in W and 1 near Y . Let

$$\tilde{h} := h|_{S_{i_1}, \dots, S_{i_l}} \circ \pi_{i_1} \circ \dots \circ \pi_{i_l}.$$

We perturb λH to

$$\tilde{H} := \lambda H + \delta_Y \rho \tilde{h}.$$

For δ_Y sufficiently small, there are no fixed points in the region where the derivative on ρ is non-zero by a compactness argument. The point is that if there were such orbits for δ_Y small, then there would be a sequence of δ_Y 's converging to zero with a fixed point in the region where $d\rho \neq 0$. But this would imply by compactness that λH would also have such an orbit which is a contradiction. There are no 1-periodic orbits of $\tilde{H}$ near the boundary of Y for the following reason: We have that Y is a manifold with corners. Near a codimension $k \geq 1$ corner of Y ,

$$H = \nu(r_{i_1}) + \dots + \nu(r_{i_l}) + \nu(r_{j_1}) + \dots + \nu(r_{j_k})$$

for some $j_1 \dots j_k$ not equal to any of $i_1, \dots, i_l$. Hence in this region $\tilde{H}$ is equal to

$$\begin{aligned} & \nu(r_{i_1}) + \dots + \nu(r_{i_l}) + \nu(r_{j_1}) + \dots + \nu(r_{j_k}) + \\ & \delta_Y (\nu(r_{i_1} - \delta_h) + \dots + \nu(r_{i_l} - \delta_h) + \nu(r_{j_1}(1 - \delta_h)) + \dots + \nu(r_{j_k}(1 - \delta_h))). \end{aligned}$$

Let $J := \{i_1, \dots, i_l, j_1, \dots, j_k\}$. Near this codimension k corner, the Hamiltonian $\tilde{H}$ preserves the fibers $\prod_{j \in J} \mathbb{D}_\epsilon$ of:

$$\pi_{j_1} \circ \dots \circ \pi_{j_k} \circ \pi_{i_1} \circ \dots \circ \pi_{i_l}.$$

These Hamiltonians also split up as a sum:

$$\sum_{j \in J} \nu(r_j) + \delta_Y(\nu(r_j(1 - \delta_h)))$$

on this fiber $\prod_{j \in J} \mathbb{D}_\epsilon$. Near the boundary of the closure of $\nu(r_j) > 0$, $\nu(r_j) + \delta_Y(\nu(r_j(1 - \delta_h)))$ has no 1-periodic orbits for δ_Y, δ_h sufficiently small. Hence $\tilde{H}$ has no 1-periodic orbits near this codimension k corner of Y . This implies that $\tilde{H}$ has no 1-periodic orbits on a small neighbourhood of ∂Y .

Hence all the fixed points of $\tilde{H}$ are contained in the region where $\nu(r_k) = 0$ for all such k satisfying $k \neq i_j \forall j$. In this region, we have that the Hamiltonians $\tilde{h}$ and λH Poisson commute because the Hamiltonian vector field of $\tilde{h}$ is contained in the horizontal plane distribution of the fibration $\pi_{i_1} \circ \dots \circ \pi_{i_l}$ which is contained in the level set of λH . Let $p \in Y$ be a fixed point of $\tilde{H}$. The symplectic tangent bundle splits into $\mathbb{R}^{2l} \oplus \mathbb{R}^{2n-2l}$. Here $\mathbb{R}^{2l}$ is the tangent space of the fiber of $\pi_{i_1} \circ \dots \circ \pi_{i_j}$ and $\mathbb{R}^{2n-2l}$ is the symplectic complement (this is the Horizontal plane bundle coming from the natural connection on this fibration). The linearized return map restricted to $\mathbb{R}^{2l}$ is the same as the linearized return map of the Hamiltonian $\sum_{j=1}^l \nu(r_{i_j})$ restricted one of the fibers of $\pi_{i_1} \circ \dots \circ \pi_{i_j}$. Because this fiber is symplectomorphic to a product of disks, we get that the flow is equal to the flow of $\sum_{j=1}^l \nu(R_j)$ on $(\mathbb{D}_\epsilon)^l$ where R_j is the j 'th radial coordinate on $\mathbb{D}_\epsilon$. Because this Hamiltonian system splits as a product of autonomous non-degenerate Hamiltonians, we have that this Linearized return map has an l dimensional eigenspace with eigenvalue 1. This eigenspace is spanned by the Hamiltonian vectors $X_{\nu(R_j)}$ $j = 1, \dots, l$ at this point. The linearized return map of $\tilde{h}$ when restricted to the symplectic complement $\mathbb{R}^{2n-2l}$ is conjugate to the linearized return map of the Hamiltonian $\delta_Y h|_{\cap_j S_{i_j}}$ inside $\cap_j S_{i_j}$. Because this is a C^2 small Morse function (for $\delta_Y > 0$ sufficiently small), this has no eigenvalue equal to 1. This implies that the linearized return map at p of the autonomous Hamiltonian $\tilde{H}$ has a $2l$ dimensional eigenspace with eigenvalue 1 which is spanned by the Hamiltonian vectors $X_{\nu(R_j)}$ at p . Also the number of critical points created from the manifold Y of fixed points is at most the number of critical points of $h|_{\cap_j S_{i_j}}$. If we perturb all such manifolds of fixed points of λH we get an non-degenerate autonomous Hamiltonian with at most $C\lambda^d (S^1)^l$ families of fixed points where C is some constant greater than the sum over all strata $\cap_j S_{i_j}$ of the number of critical points of $h|_{\cap_j S_{i_j}}$ multiplied by $2\pi \cdot 2^{nw}$. We will also perturb the fixed points where $\lambda H = 0$ so that it becomes non-degenerate in this region. Let M_H be the number of such fixed points (such a number can be independent of λ).

Finally using work from [CFHW96], we perturb each $(S^1)^l$ family of orbits away from the divisors $\cup_i S_i$ into 2^l orbits creating a Hamiltonian $\varrho : S^1 \times W \rightarrow \mathbb{R}$ where all of its orbits are non-degenerate. Really [CFHW96] deals with S^1 families of periodic orbits and not $(S^1)^l$ families, but because our Hamiltonian $\tilde{H}$ restricted to each fiber $\mathbb{R}^{2l}$ splits up as a sum $\sum_j \nu(R_j)$ we can perturb $\nu(R_j)$ first (if we view such a function as a function on one of the $\mathbb{R}^2$ factors of $\mathbb{R}^{2l}$). Hence the number of orbits of ϱ is bounded above by $2^{\dim_{\mathbb{R}} A} C \lambda^d + M_H$. This implies that H is P bounded where P is the polynomial $2C \lambda^d$. $\square$

Proof. of Theorem 6.3. From the discussion before Lemma 6.8 we can find a Hamiltonian H which is compatible with a compactification of A . We can choose an appropriate compactification M of A so that the constant d in Lemma 6.8 is equal to m_A . Hence by this lemma we have that H satisfies the Liouville vector field property and is P bounded where P is a polynomial of degree at most m_A .

Because A, B are convex deformation equivalent at infinity, by Lemma 6.7 we have that B also admits a Hamiltonian H' that satisfies the Liouville vector field property and is P bounded. By Lemma 6.6 we then have that $(SH_*^\#(B, \theta_B))$ is isomorphic to a filtered directed system (V_λ) where $|V_\lambda| \leq P(\lambda)$. This proves the Theorem. $\square$

7. FINITE COVERS OF SMOOTH AFFINE VARIETIES

In this section we will prove Theorem 1.5. We will prove the following which has Theorem 1.5 as a direct corollary:

Theorem 7.1. *Suppose that a finite type convex symplectic manifold B and a finite cover $\tilde{A}$ of a smooth affine variety A are convex deformation equivalent at infinity. Then filtered directed system $(SH_*^\#(B, \theta_B, \lambda))$ is isomorphic to a filtered directed system (V_x) where the dimension $|V_x|$ satisfies $|V_x| \leq Q(x)$ for some polynomial Q . The degree of this polynomial is less than or equal to m_A where m_A is defined in the introduction.*

Proof. of Theorem 7.1. By looking at the proof of Theorem 6.3 we see that A admits a Hamiltonian H which satisfies the Liouville vector field property and which is P bounded for some polynomial P of degree m_A .

Let $p : \tilde{A} \rightarrow A$ be the covering of degree k . We can pull back this Hamiltonian H to $\tilde{H}$. This also satisfies the Liouville vector field property and is also kP bounded. It satisfies the Liouville vector field property because we can lift any Liouville vector field on A to a Liouville vector field on $\tilde{A}$. It is kP bounded for the following reason: If H' is any Hamiltonian and $\tilde{H}'$ its lift to $\tilde{A}$ then any orbit of $\tilde{H}'$ projects down to an orbit of H' . This orbit is non-degenerate if and only if the projected orbit is non-degenerate. There are at most k orbits of $\tilde{H}'$ that project to a given orbit of H . Hence the number of non-degenerate orbits of $\tilde{H}'$ is bounded above by k times the

number of non-degenerate orbits of H' . Also the action of this orbit is equal to the action of the projected orbit.

Hence by Lemma 6.7, B admits a Hamiltonian which is kP bounded. Hence by Lemma 6.6 we get that $(SH_*^\#(B, \theta_B, \lambda))$ is isomorphic to (V_x) where the dimension $|V_x|$ satisfies $|V_x| \leq kP(x)$. Hence we have proven this theorem where $Q = kP$. $\square$

8. APPENDIX A: CONVEX SYMPLECTIC MANIFOLDS

Recall that a convex symplectic manifold is a manifold M together with a 1-form θ

- (1) $\omega := d\theta$ is a symplectic form.
- (2) The ω -dual X_θ of θ is a vector field satisfying $df_M(X_\theta) > 0$ in the region $f_M^{-1}(A_M)$ where $A_M \subset \mathbb{R}$ is an unbounded subset and $f_M : M \rightarrow \mathbb{R}$ an exhausting function.

Exhausting means that f_M is bounded from below and the preimage of every compact set is compact.

We now need to describe various ways of deforming (M, θ) . The first kind of deformation is the most restrictive: Two convex symplectic manifolds $(M, \theta), (M', \theta')$ are *strongly deformation equivalent* if there is a diffeomorphism $\phi : M \rightarrow M'$, an exhausting function $g : M \rightarrow \mathbb{R}$ and a smooth family of 1-forms θ_t ($t \in [0, 1]$) on M such that:

- (1) $\omega_t := d\theta_t$ is symplectic.
- (2) If X_{θ_t} is the ω_t -dual of θ_t then $dg(X_{\theta_t}) > 0$ on $g^{-1}(A_g)$ where $A_g \subset [0, \infty)$ is unbounded.
- (3) $\theta_0 = \theta$ and $\theta_1 = \phi^*\theta'$.

Two convex symplectic manifolds $(M, \theta), (M', \theta')$ are *convex deformation equivalent* if there is a finite sequence $(M_1, \theta_1), \dots, (M_k, \theta_k)$ such that

$$(M_1, \theta_1) = (M, \theta), (M_k, \theta_k) = (M', \theta')$$

and such that for each $i < k$, (M_i, θ_i) is strongly deformation equivalent to (M_{i+1}, θ_{i+1}) .

An important class of convex symplectic manifolds are the complete ones. A *complete convex symplectic manifold* is a convex symplectic manifold such that the associated Liouville vector field V (the ω -dual of θ) is integrable for all time. A (strong) convex deformation (M, θ_t) is called *complete* if the associated Liouville vector fields X_{θ_t} are integrable.

Lemma 8.1. *Let (M, θ) be a convex symplectic manifold and let $K \subset M$ be any compact set. There exists a smooth family of 1-forms $\theta_t, (t \in [0, 1])$ on M such that:*

- (1) $\theta_0 = \theta$
- (2) $\omega_t := d\theta_t$ is a symplectic form.
- (3) *If X_{θ_t} is the ω_t -dual of θ_t , then there is a smooth family of functions $f_t : M \rightarrow \mathbb{R}$ such that $X_{\theta_t} = f_t X_{\theta_0}$ and $0 < f_t \leq 2$.*

- (4) The vector field X_{θ_1} is integrable.
- (5) $\theta_t|_K = \theta_0|_K$.

Proof. of Lemma 8.1. Let X_θ be the $d\theta$ -dual of θ . Let $A \subset [0, \infty)$ be an unbounded subset and f an exhausting function such that $df(X_\theta) > 0$ on $f^{-1}(A)$. Note that each $a \in A$ is a regular value of f because $df(X_\theta) > 0$ on $f^{-1}(a)$ which implies in particular that $df \neq 0$ on $f^{-1}(a)$. Because f is smooth, there exists a sequence $c_1 < c_2, \dots$ tending to infinity such that c_i is in A and X_θ is transverse to $f^{-1}(c_i)$ and pointing outwards. We choose c_i large enough so that $f^{-1}(-\infty, c_i)$ contains K . By flowing along X_θ there is a neighbourhood of the manifold $C_i := f^{-1}(c_i)$ diffeomorphic to $[1 - \epsilon_i, 1 + \epsilon_i] \times C_i$ with $\theta = r_i \alpha_i$ where r_i parameterizes the interval and $\alpha_i = \theta|_{C_i}$. Again we make these neighborhoods small enough so that they are disjoint from K . Let $\mathcal{N}_i$ be this neighbourhood and let $\mathcal{N}$ be the union of all these neighborhoods. We can make these neighborhoods small enough so that they are disjoint and also so that $\epsilon_i \leq \frac{1}{2}$.

We will now construct θ_t : Let $g_i : [1 - \epsilon_i, 1 + \epsilon_i] \rightarrow \mathbb{R}$ be smooth functions such that:

- (1) $g'_i > 0$.
- (2) In the region $[1 - \epsilon_i, 1 - 2\epsilon_i/3]$, we have $g_i(r_i) = r_i$.
- (3) In the region $[1 - \epsilon_i/2, 1 + \epsilon_i/2]$ we have $g_i/g'_i \leq \epsilon_i$.
- (4) $g_i/g'_i \leq 1$ or equivalently $\log(g_i)' \geq 1$.
- (5) In the region $[1 + 2\epsilon_i/3, 1 + \epsilon_i]$ we have $g_i(r_i) = \kappa_i r_i$ where $\kappa_i \geq 1$ is some large constant.

We define $g_i^t(r_i) := (1 - t)r_i + tg_i(r_i)$. We define $c_0 := -\infty$. Let $\Xi_i := \prod_{j=1}^{i-1} \kappa_j$ for $i > 1$ and $\Xi_1 := 1$. In the region $f^{-1}(c_{i-1}, c_i) \setminus \mathcal{N}$, we define $\theta_t := ((1 - t) + t\Xi_i)\theta$. In the region $\mathcal{N}_i$ we define θ_t as $((1 - t) + t\Xi_i)g_i^t(r_i)\alpha_i$. This definition ensures that θ_t ($t \in [0, 1]$) is a smooth family of 1-forms.

Outside $\mathcal{N}_i$, we have that θ_t is equal to some locally constant function multiplied by θ , hence $d\theta_t$ is still symplectic. In the region $\mathcal{N}_i$, we have that $d\theta_t = ((1 - t) + t\Xi_i)((g_i^t)'dr_i \wedge \alpha_i + r_i d\alpha_i)$. This is symplectic because α_i is a contact form and $g_i^{t'} > 0$. Hence $d\theta_t$ is symplectic for all t . Outside $\mathcal{N}_i$ we have that the ω_t -dual X_{θ_t} of θ_t is equal to V because rescaling θ by a locally constant function does not change the associated dual vector field. Inside $\mathcal{N}_i$ we have that $X_{\theta_t} = g_i^t/g_i^{t'} \frac{\partial}{\partial r_i}$ which is equal to $V = r_i \frac{\partial}{\partial r_i}$ multiplied by some positive function f_t . Because $g_i/g'_i \leq \epsilon_i$ in the region $[1 - \epsilon_i/2, 1 + \epsilon_i/2]$, if we flow any point p in the region where $r_i = 1 - \epsilon_i/2$ along the vector field X_{θ_1} for time 1 to a point q , then q is still contained in $[1 - \epsilon_i/2, 1 + \epsilon_i/2] \times C_i$. This ensures that the vector field X_{θ_1} is complete and hence (M, θ_1) is complete. We also have that $f_t \leq 2$ because $f_t = 1$ outside $\mathcal{N}$ and is equal to $\frac{g_i^t}{g_i^{t'} r_i} \leq \frac{1}{1 - \epsilon_i} \leq 2$ inside $\mathcal{N}_i$. Finally we have that $\theta_0 = \theta$ by definition and $\theta_t|_K = \theta_0|_K$ because $\mathcal{N}$ is disjoint from K . $\square$

We also have a 1-parameter version of this Lemma as follows (with almost exactly the same proof): Suppose (M, θ_s) ($s \in [0, 1]$) is a smooth family of

convex symplectic manifolds such that we have a function $f : M \rightarrow \mathbb{R}$ and an unbounded $A \subset [0, \infty)$ with $df(X_{\theta_s}) > 0$ on $f^{-1}(A)$ where X_{θ_s} is the $d\theta_s$ -dual of θ_s . Then there is a two parameter family of 1-forms $\theta_{s,t}$ ($s, t \in [0, 1]^2$) satisfying:

- (1) $\theta_{s,0} = \theta_s$
- (2) $\omega_{s,t} := d\theta_{s,t}$ is a symplectic form.
- (3) If $X_{\theta_{s,t}}$ is the $\omega_{s,t}$ -dual of $\theta_{s,t}$, then there is a smooth family of functions $f_{s,t}$ such that $X_{\theta_{s,t}} = f_{s,t}X_{\theta_{s,0}}$ and $0 < f_{s,t} \leq 2$.
- (4) The vector field $X_{\theta_{s,1}}$ is integrable for each $s \in [0, 1]$.

Corollary 8.2. *Every convex symplectic manifold is strongly deformation equivalent to a complete convex symplectic manifold.*

Proof. Let f_M, X_θ be the associated function and Liouville vector field of M . There is an unbounded $A \subset [0, \infty)$ such that $df_M(X_\theta) > 0$ in $f^{-1}(A)$. Let (M, θ_t) be the family of convex symplectic manifolds described in the Lemma 8.1 above. If X_{θ_t} is the associated Liouville vector field of (M, θ_t) then because $X_{\theta_t} = f_t X_\theta$ for some smooth family of functions $f_t > 0$, we have that $df_M(X_{\theta_t}) > 0$ in $f^{-1}(A)$. Hence we have that (M, θ_t) is our strong convex deformation. $\square$

Corollary 8.3. *Let f_M, X_θ be the associated function and Liouville vector field of M . Suppose M is of finite type which means that $df_M(X_\theta) > 0$ in the region $f^{-1}[C, \infty)$, then M is convex deformation equivalent to the completion of the Liouville domain $f^{-1}(-\infty, C]$.*

Proof. Let (M, θ_t) be the family of 1-forms as in Lemma 8.1 and let $X_{\theta_t} = f_t X_\theta$ be the associated family of Liouville vector fields. Then $df_M(X_{\theta_1}) > 0$ in the region $f_M^{-1}[C, \infty)$. Let $D := f_M^{-1}(-\infty, C]$ and let $\alpha = \theta_1|_{\partial D}$. By flowing the contact manifold $(\partial D, \alpha)$ along X_{θ_1} (which is integrable) we obtain a diffeomorphism ϕ from $\partial D \times [1, \infty)$ to $f_M^{-1}[C, \infty)$ such that $\phi^* \theta_1 = r\alpha$ where r parameterizes $[1, \infty)$. Also by the previous corollary 8.2 we have that (M, θ) is convex deformation equivalent to (M, θ_1) . Hence (M, θ_1) is convex deformation equivalent to the completion $(\widehat{D}, \theta_1)$. $\square$

Suppose we have a convex deformation equivalence (M, θ_t) . We say that this is a *complete convex deformation equivalence* if the $d\theta_t$ -dual X_{θ_t} of θ_t is an integrable vector field for each $t \in [0, 1]$.

Lemma 8.4. *Let $(M, \theta), (M', \theta')$ be complete convex symplectic manifolds that are convex deformation equivalent. Then there is a complete convex deformation equivalence between (M, θ) and (M', θ') .*

We need a preliminary Lemma first:

Lemma 8.5. *Suppose $(M, \theta), (M', \theta')$ are complete convex symplectic manifolds that are strongly deformation equivalent. Then there is a complete strong deformation equivalence between them.*

Proof. of Lemma 8.5. Let (M, θ_s) be the strong deformation equivalence. Let X_{θ_s} be the associated Liouville vector fields and let $f : M \rightarrow \mathbb{R}$ and $A \subset [0, \infty)$ (unbounded) be such that $df(X_{\theta_s}) > 0$ on $f^{-1}(A)$. By Lemma 8.1 there is a two parameter family of 1-forms $\theta_{s,t}((s, t) \in [0, 1]^2)$ such that:

- (1) $\theta_{s,0} = \theta_s$
- (2) $\omega_{s,t} := d\theta_{s,t}$ is a symplectic form.
- (3) If $X_{\theta_{s,t}}$ is the $\omega_{s,t}$ -dual of $\theta_{s,t}$, then there is a smooth family of functions $f_{s,t}$ such that $X_{\theta_{s,t}} = f_{s,t}X_{\theta_{s,0}}$ and $0 < f_{s,t} \leq 2$.
- (4) The vector field $X_{\theta_{s,1}}$ is integrable for each $s \in [0, 1]$.

Let $p : [0, 1] \rightarrow [0, 1]^2$ be a smooth path starting at $(0, 0)$ and ending at $(1, 0)$ whose image is equal to $\{0\} \times [0, 1] \cup [0, 1] \times \{1\} \cup \{1\} \times [0, 1]$. Then $(M, \theta_{p(t)})$ is our complete strong convex deformation. This is because $X_{\theta_{s,0}}$ (resp. $X_{\theta_{s,1}}$) is integrable because it is equal to $f_{s,0}X_{\theta_{0,0}}$ (resp. $f_{s,1}X_{\theta_{0,1}}$) with $0 < f_{s,t} \leq 2$ for all s, t . This is also because $f_{1,t}$ is integrable for all t . This completes the Lemma. $\square$

Proof. of Lemma 8.4. Let (M, θ_s) be the convex deformation between (M, θ) and (M', θ') . This means that we break up (M, θ_s) into a finite number of strong deformations $(M, \theta_s^1), \dots, (M, \theta_s^k)$. Let V_s^i be the associated Liouville vector fields. For a given strong deformation (M, θ_s^i) there is an unbounded set $A_i \subset [0, \infty)$ and a function $g_i : M \rightarrow \mathbb{R}$ such that $g_i(V_s^i) > 0$ in $g_i^{-1}(A_i)$. By the parameterized version of Lemma 8.1 we can replace θ_s^i with $\theta_{s,t}^i$ such that:

- (1) $\theta_{s,0}^i = \theta_s^i$
- (2) $\omega_{s,t}^i := d\theta_{s,t}^i$ is a symplectic form.
- (3) If $V_{s,t}^i$ is the $\omega_{s,t}^i$ -dual of $\theta_{s,t}^i$, then there is a smooth family of functions $f_{s,t}^i$ such that $V_{s,t}^i = f_{s,t}^i V_{s,0}^i$ and $0 < f_{s,t}^i \leq 2$.
- (4) The vector field $V_{s,1}^i$ is integrable for each $s \in [0, 1]$.

We now replace the deformation (M, θ_s^1) by the concatenation $(M, \theta_s'^1)$ of the homotopies: $(M, \theta_{0,t}^1)$ and $(M, \theta_{t,1}^1)$. This is still a strong deformation because $g_1(V_{s,t}^1) > 0$ in the region $g_1^{-1}(A_1)$. For $1 < i < k$ we replace (M, θ_s^i) with the concatenation $(M, \theta_s'^i)$ of: $(M, \theta_{1,1-t}^{i-1})$, $(M, \theta_{0,t}^i)$ and $(M, \theta_{t,1}^i)$. This again is a strong deformation using the function g_i and unbounded set $A_i \subset [0, \infty)$. The reason why $g_i(V_{1,1-t}^{i-1}) > 0$ in $g_i^{-1}(A_i)$ is because $V_{1,1-t}^{i-1}$ is $f_{1,1-t}^{i-1} V_t^{i-1}$ where $f_{1,1-t}^{i-1} > 0$. Finally we replace the deformation (M, θ_s^k) with the concatenation $(M, \theta_s'^k)$ of $(M, \theta_{1,1-t}^{k-1})$, $(M, \theta_{0,t}^k)$, $(M, \theta_{t,1}^k)$ and $(M, \theta_{1,1-t}^k)$. Hence we have a new convex deformation equivalence $(M, \theta_t'^1), \dots, (M, \theta_t'^k)$ with the property that θ_0^i, θ_1^i is complete for all i . By Lemma 8.5, we can replace this with a new convex deformation equivalence: $(M, \theta_t''^1), \dots, (M, \theta_t''^k)$ with the property that $\theta_t''^i$ is complete for all $1 \leq i \leq k, t \in [0, 1]$. This is a complete convex deformation equivalence. $\square$

Corollary 8.6. *Suppose that $(M, \theta), (M, \theta')$ are complete convex symplectic manifolds that are convex deformation equivalent. Then they are exact symplectomorphic to each other.*

Proof. of 8.6. By Lemma 8.4 there is a complete convex deformation equivalence between (M, θ) and (M, θ') . Hence by [CE, Proposition 12.2] we get that they are exact symplectomorphic. $\square$

We also need another lemma similar to corollary 8.6 except that we will be dealing with convex deformation equivalence at infinity as in definition 6.2.

Lemma 8.7. *Suppose that $(W, \theta_W), (W', \theta_{W'})$ are complete finite type convex symplectic manifolds that are convex deformation equivalent at infinity. Then they are exact symplectomorphic at infinity.*

Proof. of Lemma 8.7. By the definition of convex deformation equivalent at infinity, we have a sequence of convex symplectic manifolds: (Q_i, θ_{Q_i}) for $i = 1, \dots, k$ such that:

- (1) Q_1 is convex deformation equivalent to W_1 .
- (2) Q_k is convex deformation equivalent to W_2 .
- (3) Q_i is convex deformation equivalent to Q_{i+1} or they are exact symplectomorphic at infinity.

First of all (by using the fact that the identity map is an exact symplectomorphism and also a convex deformation equivalence), we can assume that k is even and that for odd i , that Q_i, Q_{i+1} are convex deformation equivalent and for even i that Q_i, Q_{i+1} are exact symplectomorphic at infinity.

Suppose we have two convex symplectic manifolds $(A, \theta_A), (B, \theta_B)$ that are exact symplectomorphic at infinity. We wish to find two complete convex symplectic manifolds $\tilde{A}$ and $\tilde{B}$ that are exact symplectomorphic at infinity and such that $\tilde{A}$ is convex deformation equivalent to A and $\tilde{B}$ is convex deformation equivalent to B . Let $\phi : A \setminus K_A \rightarrow B \setminus K_B$ be the exact symplectomorphism at infinity where K_A, K_B are relatively compact sets. By possibly enlarging K_A a little bit, we can ensure that there is a function $f : A \rightarrow \mathbb{R}$ such that $\phi^* \theta_B = \theta_A + df$. Let $\mathcal{N}$ be a small neighbourhood of the closure of K_A . By Lemma 8.1, there is a family of 1-forms θ_A^s such that

- (1) $d\theta_A^s$ is symplectic.
- (2) $\theta_A + df = \theta_A^0$.
- (3) $\theta_A^s|_{\mathcal{N}} = \theta_A^0|_{\mathcal{N}}$.
- (4) θ_A^1 is complete.
- (5) The $d\theta_A^s$ -dual $X_{\theta_A^s}$ of θ_A^s satisfies $X_{\theta_A^s} = gX_{\theta_A^0}$ for some positive function g .

We can also define θ_B^s to be equal to θ_B near K_B and equal to $\phi_*(\theta_A^s)$ outside K_B .

These are all convex symplectic manifolds because their Liouville vector fields are all of the form $g_s X_{\theta_A + df}$ (resp. $g_s X_{\theta_B}$) for some family of functions g_s and the same reason ensures that this is a convex deformation. Also by

[McL09, Lemma 8.3], we have that (A, θ_A) is convex deformation equivalent to $(A, \theta_A + df)$. This means that (A, θ_A) (resp. (B, θ_B)) is convex deformation equivalent to the complete convex symplectic manifold $\tilde{A} := (A, \theta_A^1)$ (resp. $\tilde{B} := (B, \theta_B^1)$). Also $\tilde{A}, \tilde{B}$ are exact symplectomorphic at infinity.

The previous discussion ensures (by changing the convex deformation equivalences from Q_i to Q_{i+1}) that we can assume that the convex symplectic manifolds Q_i are all complete convex symplectic manifolds. Note we can also assume that Q_1 and Q_k are complete because W and W' are complete. By Corollary 8.6 this implies that for all odd i , Q_i is exact symplectomorphic to Q_{i+1} and hence in particular they are exact symplectomorphic at infinity. Because the property of being exact symplectomorphic at infinity is transitive, we have that W, W' are exact symplectomorphic at infinity. This proves the Lemma. $\square$

9. APPENDIX B: A MAXIMUM PRINCIPLE

Let N be a manifold and θ a 1-form on N so that

- (1) $\omega := d\theta$ is a symplectic form.
- (2) The ω -dual X_θ of θ is transverse to the boundary of N and pointing inwards.

Let S be a compact Riemann surface with boundary and complex structure j and γ a 1-form on S . Let $H : S \times N \rightarrow \mathbb{R}$ be a family of Hamiltonians parameterized by S . We sometimes write this as a family of functions $H_\sigma : N \rightarrow \mathbb{R}$ parameterized by $\sigma \in S$. Let J_σ be a family ω compatible almost complex structures parameterized by $\sigma \in S$. A small neighbourhood of ∂N is diffeomorphic to $\partial N \times [1, 1 + \epsilon_N)$ where $\theta = r\alpha$. Here r parameterizes the interval $[1, 1 + \epsilon_N)$ and $\alpha = \theta|_{\partial N}$. We require that $\theta \circ J_\sigma = dr$ and that $H_\sigma = f(r)$ for some function f with $f(1) = 1$ and $f'(1) = 1$ near ∂N . The differential dH uniquely splits up as $d_S H + d_N H$ where vectors tangent to N are contained in the kernel of $d_S H$ and vectors tangent to S are in the kernel of $d_N H$. We can view $d_S H$ as a family of 1-forms on S parameterized by N , so for each $p \in N$ we define $d_S H_{(\cdot, p)}$ to be $d_S H$ restricted to $S \times \{p\}$. We require that $d_S H_{(\cdot, p)} \wedge \gamma + H(\cdot, p)d(\gamma) \geq 0$ for each $p \in N$.

Let $u : S \rightarrow N$ satisfy: $(du - X_{H_\sigma} \otimes \gamma)_{J_\sigma}^{0,1} = 0$ at each point $\sigma \in S$. In other words:

$$(3) \quad du - X_{H_\sigma} \otimes \gamma + J_\sigma \circ (du - X_{H_\sigma} \otimes \gamma) \circ j = 0.$$

The aim of this section is to prove:

Lemma 9.1. *If $u(\partial S) \subset \partial N$ then $u(S) \subset \partial N$.*

This lemma is similar to [AS10, Lemma 7.2]. Before we prove Lemma 9.1 we need to define the *geometric energy* and *topological energy* of u . The geometric energy is defined as:

$$E^{\text{geom}} := \int_S \|du - X_{H_\sigma} \otimes \gamma\|_{J_\sigma}^2$$

where $\|\cdot\|_{J_\sigma}$ is the norm coming from the metric $\omega(\cdot, J_\sigma(\cdot))$ and some compatible metric on (S, j) . The topological energy is defined as:

$$E^{\text{top}} := \int_S u^* \omega + d(u^* H_\sigma \gamma).$$

Here $u^* H_\sigma$ is the function sending $\sigma \in S$ to $H(\sigma, u(\sigma))$. We define $u^* d_N H$ to be the 1-form on S such that for each vector V on S , $(u^* d_N H)(V) = d_N H(V, u_*(V))$. Similarly we define $(u^* d_S H)(V)$ to be $(d_S H)(V, u_*(V))$.

Let σ be a point on S and $s+it$ a local holomorphic chart around σ where ∂_s, ∂_t have magnitude 1 at $(s, t) = (0, 0)$ and where σ is the point $(0, 0)$. We have:

$$\begin{aligned} \|du - X_{H_\sigma} \otimes \gamma\|_{J_\sigma}^2 &= \omega(\partial_s u - X_{H_\sigma} \gamma(\partial_s), \partial_t u - X_{H_\sigma} \gamma(\partial_t)) \\ &= u^* \omega(\partial_s, \partial_t) - \gamma(\partial_s) d_N H(\partial_t u) + \gamma(\partial_t) d_N H(\partial_s u) \\ &= u^* \omega(\partial_s, \partial_t) + (u^* (d_N H) \wedge \gamma)(\partial_s, \partial_t). \end{aligned}$$

Also $d_N H(\partial_t u)$ means that we consider $\partial_t u$ as a vector inside $\{\sigma\} \times N$ and then contract it with dH . The expression $d_N H(\partial_s u)$ has a similar meaning. Hence $E^{\text{geom}}(u) = \int_S u^* \omega + \int_S u^* d_N H \wedge \gamma$. We have that

$$d(u^* H_\sigma \gamma) = u^* d_N H \wedge \gamma + u^* d_S H \wedge \gamma + u^* H_\sigma d\gamma.$$

Hence we have:

$$E^{\text{geom}}(u) = E^{\text{top}}(u) - \int_S (u^* d_S H \wedge \gamma + u^* H_\sigma d\gamma)$$

which implies that $E^{\text{geom}}(u) \leq E^{\text{top}}(u)$.

Proof. of Lemma 9.1 We suppose that $u(\partial S) \subset \partial N$. We just need to show that $E^{\text{geom}}(u) = 0$ as this will force our surface S to map to a Reeb orbit of ∂N . We have that $E^{\text{top}}(u) \geq E^{\text{geom}}(u)$, so we now need to show that $E^{\text{top}}(u) = 0$. By Stokes' theorem we have:

$$E^{\text{top}}(u) = \int_{\partial S} u^* \theta + u^* H_\sigma \gamma.$$

Because $H_\sigma = f(r)$ with $f'(1) = 1$, we have that $-X_{H_\sigma}$ is equal to the Reeb vector field on ∂N . Hence $\theta(X_{H_\sigma}) = -1 = -H_\sigma$ along ∂N as $f(1) = 1$. So

$$\begin{aligned} E^{\text{top}}(u) &= \int_{\partial S} \theta \circ (du - X_{H_\sigma} \otimes \gamma) \\ &= \int_{\partial S} \theta \circ J_\sigma \circ (du - X_{H_\sigma} \otimes \gamma) \circ (-j) = \int_{\partial S} dr \circ (du - X_{H_\sigma} \otimes \gamma) \circ (-j). \end{aligned}$$

Because $dr(X_{H_\sigma}) = 0$ our integral becomes:

$$E^{\text{top}}(u) = - \int_{\partial S} dr \circ du \circ j.$$

If a vector V is tangent to ∂S and pointing in the direction in which ∂S is oriented then $j(V)$ points inwards. This implies that $dr \circ du \circ j(V) \geq 0$ because r increases as we move towards the interior of N . Hence

$$E^{\text{top}}(u) \leq 0.$$

Hence $E^{\text{geom}}(u)$ vanishes which gives us our result. $\square$

Here are two applications of this lemma: Let M be a Liouville domain with 1-form θ_M . Then its completion $\widehat{M}$ has a cylindrical end $\partial M \times [1, \infty)$ with cylindrical coordinate r_M . We define $K : \widehat{M} \rightarrow \mathbb{R}$ to be an autonomous Hamiltonian on $\widehat{M}$ which is equal $k(r_M)$ near ∂M where $k' > 0$. Let $H_{s,t}$ be a family of Hamiltonians parameterized by $(s, t) \in \mathbb{R} \times S^1$ so that $H_{s,t} = K + a_s$ near ∂M where a_s is a smooth family of constants. We require that $\frac{\partial H}{\partial s} \geq \frac{\partial a_s}{\partial s}$. We define $J_{s,t}$ to be a smooth family of almost complex structures which are cylindrical near ∂M .

Corollary 9.2. *Suppose that $u_1 : \mathbb{R} \times S^1 \rightarrow \widehat{M}$ satisfies the perturbed Cauchy-Riemann equations:*

$$\partial_s u_1 + J_{s,t} \partial_t u_1 = J_{s,t} X_{H_{s,t}}$$

and that $u_1(s, t) \in M$ for $|s| \gg 1$. Then $u_1(s, t) \in M$ for all $(s, t) \in \mathbb{R} \times S^1$.

Proof. of Corollary 9.2. Let S be equal to $u_1^{-1}(\partial M \times [1, \infty))$. We perturb ∂M slightly so that S is a codimension 0 submanifold with boundary. We define $H_1 := \frac{H - k(1) - a_s}{k'(1)} + 1$. We let $\gamma = k'(1)dt$ and so $(d_S H_1)_{(\cdot, p)} \wedge \gamma + (H_1)_{(\cdot, p)} d\gamma \geq 0$. We have that u_1 satisfies the perturbed Cauchy Riemann equations with respect to H_1 and $J_{s,t}$. We also have that H_1 is equal to $h(r_M) = \frac{k(r_M) - k(1)}{k'(1)} + 1$ near ∂M and so $h(1) = h'(1) = 1$. By using Lemma 9.1 with $N = \partial M \times [1, \infty)$ we have that $u_1(S)$ must be contained inside ∂M . Hence the image of u_1 is contained in M . $\square$

Corollary 9.3. *Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a function satisfying $g'(s) \geq 0$. Suppose in addition that $H \geq k(1) - k'(1) + a_s$ inside $\partial M \times [1, \infty)$, then any solution: $u_2 : \mathbb{R} \times S^1 \rightarrow \widehat{M}$ of*

$$\partial_s u_2 + J_{s,t} \partial_t u_2 = J_{s,t} X_{g(s)H_{s,t}}$$

with $u_2(s, t) \in M$ for $|s| \gg 1$ has image contained in M .

Proof. of 9.3. Again we define $H_1 := \frac{H - k(1) - a_s}{k'(1)} + 1$. We let $\gamma = k'(1)g(s)dt$. Because $H_1, \frac{\partial H_1}{\partial s} \geq 0$, we have $(d_S H_1)_{(\cdot, p)} \wedge \gamma + (H_1)_{(\cdot, p)} d\gamma \geq 0$. We then apply Lemma 9.1 to give us our result. $\square$

REFERENCES

- [Abo10] Abouzaid, Mohammed. A cotangent fibre generates the Fukaya category. pages 1–40, 2010, arXiv:1003.4449.
- [Ano81] D. V. Anosov. Some homotopies in a space of closed curves. *Math. USSR-Izv.* 17:423–453, 1981.

- [AS06] Alberto Abbondandolo and Matthias Schwarz. On the Floer homology of cotangent bundles. *Comm. Pure Appl. Math.*, 59(2):254–316, 2006.
- [AS10] Mohammed Abouzaid and Paul Seidel. An open string analogue of Viterbo functoriality. *Geom. Topol.*, 14(2):627–718, 2010.
- [BEH⁺03] F. Bourgeois, Y. Eliashberg, H. Hofer, K. Wysocki, and E. Zehnder. Compactness results in symplectic field theory. *Geom. Topol.*, 7:799–888, 2003, arXiv:SG/0308183.
- [BTE09] F. Bourgeois and Yakov Eliashberg Tobias Ekholm. Effect of legendrian surgery. pages 1–79, 2009, arXiv:SG/0911.0026.
- [CE] K. Cieliebak and Y. Eliashberg. Symplectic geometry of Stein manifolds. *In preparation*.
- [CFHW96] K. Cieliebak, A. Floer, H. Hofer, and K. Wysocki. Applications of symplectic homology II: stability of the action spectrum. *Math. Z.*, 223:27–45, 1996.
- [DS94] S. Dostoglou and D. Salamon. Self-dual instantons and holomorphic curves. *Ann. of Math. (2)*, 139:581–640, 1994.
- [EG91] Y. Eliashberg and M. Gromov. Convex symplectic manifolds. In *Several complex variables and complex geometry Part 2*, volume 52, pages 135–162. Amer. Math. Soc., Providence, RI., 1991.
- [FHS95] A. Floer, H. Hofer, and D. Salamon. Transversality in elliptic Morse theory for the symplectic action. *Duke Math. J.*, 80:251–292, 1995.
- [FHT82] Yves Félix, Stephen Halperin, and Jean-Claude Thomas. The homotopy Lie algebra for finite complexes. *Inst. Hautes Études Sci. Publ. Math.*, (56):179–202 (1983), 1982.
- [FHT01] Yves Félix, Stephen Halperin, and Jean-Claude Thomas. *Rational homotopy theory*, volume 205 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2001.
- [FT82] Y. Félix and J.-C. Thomas. The radius of convergence of Poincaré series of loop spaces. *Invent. Math.*, 68(2):257–274, 1982.
- [Gro78] Mikhael Gromov. Homotopical effects of dilatation. *J. Differential Geom.*, 13(3):303–310, 1978.
- [Hir64] Heisuke Hironaka. Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II. *Ann. of Math. (2)* 79 (1964), 109–203; *ibid.* (2), 79:205–326, 1964.
- [Lam01] Pascal Lambrechts. The Betti numbers of the free loop space of a connected sum. *J. London Math. Soc. (2)*, 64(1):205–228, 2001.
- [McL09] M. McLean. Lefschetz fibrations and symplectic homology. *Geom. Topol.*, 13(4):1877–1944, 2009.
- [MS98] Dusa McDuff and Dietmar Salamon. *Introduction to symplectic topology*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1998.
- [Oan04] A. Oancea. A survey of Floer homology for manifolds with contact type boundary or symplectic homology. *Ensaïos Mat.*, 7, 2004, arXiv:SG/0403377.
- [Rit98] Alexander Ritter. Topological quantum field theory structure on symplectic cohomology. pages 1–60, 1998, arXiv:SG/1003.1781.
- [Sei08] P. Seidel. A biased view of symplectic cohomology. *Current Developments in Mathematics*, 2006:211–253, 2008.
- [SW06] D. A. Salamon and J. Weber. Floer homology and the heat flow. *Geom. Funct. Anal.*, 16(5):1050–1138, 2006.
- [Tot03] Burt Totaro. Complexifications of nonnegatively curved manifolds. *J. Eur. Math. Soc. (JEMS)*, 5(1):69–94, 2003.
- [Vit96] C. Viterbo. Functors and computations in Floer homology with applications, part II. *Preprint.*, 1996.

- [VP84] Micheline Vigué-Poirrier. Homotopie rationnelle et croissance du nombre de géodésiques fermées. *Ann. Sci. École Norm. Sup. (4)*, 17(3):413–431, 1984.